

G-dinaturality

ZORAN PETRIĆ

Matematički institut

SANU, Knez Mihailova 35, p.f. 367

email: zpetric@mi.sanu.ac.yu

February 1, 2008

Abstract

An extension of the notion of dinatural transformation is introduced in order to give a criterion for preservation of dinaturality under composition. An example of an application is given by proving that all bicartesian closed canonical transformations are dinatural. An alternative sequent system for intuitionistic propositional logic is introduced as a device, and a cut elimination procedure is established for this system.

1 Introduction

The aim of this paper is to introduce a generalization of the traditional notion of dinaturality and to give a geometrical criterion for preservation of dinaturality under composition. A certain importance is usually attached to this question (see [5], [1], [7] and [2]). It is useful to consult [10] to find about the historical perspective of the notion of naturality. The first extension of this notion towards our g-dinatural transformations was given by Eilenberg and Kelly in [5]. In the present paper we generalize the definition of dinaturality introduced by Dubuc and Street in [4]. The theory of g-dinatural transformations is here applied to bicartesian closed canonical transformations. It is proved that they are all dinatural in the sense of [4].

By a bicartesian closed category we mean a category equipped with finite products and coproducts, including initial and terminal objects, which is closed in the sense that for every object A , the functor $A \times _$ has the right adjoint $A \rightarrow _$. This category may serve as a framework for the categorial proof theory of intuitionistic propositional logic. However, despite that we are using a very traditional categorial object, our notation and definitions are a little bit unusual. This choice is forced by the technique that we intend to use here, and we believe that it is optimal.

Notation. For objects we use the schematic letters $A, B, C, \dots, A_1, \dots$, and for morphisms the schematic letters $f, g, h, \dots, f_1, \dots$. The product of A and B is denoted by $A \times B$, and the coproduct by $A + B$. We use O and I to specify the initial and the terminal object of a category. To denote that a morphism f has the source A and the target B we use the notation $f : A \vdash B$, and we say that f is of the type $A \vdash B$. Apart from the logical motivation for the symbol $\vdash$ instead of $\rightarrow$, we have another reason, which comes from our intention to write complex objects linearly: we use $A \rightarrow B$ instead of B^A for exponentiation, i.e. the image of B under the right adjoint of the functor $A \times _$. However, in Section 2, where we deal with a new notion of dinaturality, and which is, except for examples, self-contained, we use the standard symbol $\rightarrow$ for morphisms. Also, to avoid too many parentheses, we assume that the morphism operation $\circ$ binds more strongly than $\times, +, \rightarrow$; for example, we write

$g \circ f \times h$ for $(g \circ f) \times h$.

Bicartesian closed categories. A bicartesian closed category $\mathcal{B}$ satisfies:

For every triple A, B, C of objects from $\mathcal{B}$, we have the following *special morphisms* in $\mathcal{B}$

$$\begin{array}{ll}
\mathbf{1}_A : A \vdash A, & \delta_A^i : A \vdash A \times I, \\
\delta_A : A \times I \vdash A, & \mathbf{b}_{A,B,C}^\leftarrow : (A \times B) \times C \vdash A \times (B \times C), \\
\mathbf{b}_{A,B,C}^\rightarrow : A \times (B \times C) \vdash (A \times B) \times C, & \mathbf{c}_{A,B} : A \times B \vdash B \times A, \\
\mathbf{c}_A : A \vdash A \times I, & \mathbf{w}_A : A \vdash A \times A, \\
\mathbf{m}_A : A + A \vdash A, & \mathbf{l}_A : O \vdash A, \\
\mathbf{l}_{A,B}^1 : A \vdash A + B, & \mathbf{l}_{A,B}^2 : B \vdash A + B, \\
\epsilon_{A,B} : A \times (A \rightarrow B) \vdash B, & \eta_{A,B} : B \vdash A \rightarrow (A \times B),
\end{array}$$

and the following operations on morphisms:

$$\begin{array}{ll}
\frac{f : A \vdash B \quad g : B \vdash C}{g \circ f : A \vdash C}, & \frac{f : A \vdash B \quad g : C \vdash D}{f \rightarrow g : B \rightarrow C \vdash A \rightarrow D}, \\
\frac{f : A \vdash B \quad g : C \vdash D}{f \times g : A \times C \vdash B \times D}, & \frac{f : A \vdash B \quad g : C \vdash D}{f + g : A + C \vdash B + D}.
\end{array}$$

Also, the following equations must be satisfied

$$\begin{array}{ll}
(\text{cat } 1) & \mathbf{1}_B \circ f = f \circ \mathbf{1}_A = f, \\
(\text{cat } 2) & h \circ (g \circ f) = (h \circ g) \circ f, \\
(\times 1) & \mathbf{1}_A \times \mathbf{1}_B = \mathbf{1}_{A \times B}, \\
(\times 2) & (g_1 \circ g_2) \times (f_1 \circ f_2) = (g_1 \times f_1) \circ (g_2 \times f_2), \\
(\delta) & f \circ \delta_A = \delta_B \circ (f \times \mathbf{1}_I) \\
(\delta \delta^i) & \delta_A \circ \delta_A^i = \mathbf{1}_A, \quad \delta_A^i \circ \delta_A = \mathbf{1}_{A \times I}, \\
(\delta \mathbf{c}) & \delta_I \circ \mathbf{c} = \delta_I, \\
(\mathbf{b}) & ((f \times g) \times h) \circ \mathbf{b}_{A,B,C}^\rightarrow = \mathbf{b}_{D,E,F}^\rightarrow \circ (f \times (g \times h)), \\
(\mathbf{bb}) & \mathbf{b}_{A,B,C}^\rightarrow \circ \mathbf{b}_{A,B,C}^\leftarrow = \mathbf{1}_{(A \times B) \times C}, \quad \mathbf{b}_{A,B,C}^\leftarrow \circ \mathbf{b}_{A,B,C}^\rightarrow = \mathbf{1}_{A \times (B \times C)}, \\
(\mathbf{b5}) & \mathbf{b}_{A \times B, C, D}^\rightarrow \circ \mathbf{b}_{A, B \times C, D}^\rightarrow = (\mathbf{b}_{A, B, C}^\rightarrow \times \mathbf{1}_D) \circ \mathbf{b}_{A, B \times C, D}^\rightarrow \circ (\mathbf{1}_A \times \mathbf{b}_{B, C, D}^\rightarrow), \\
(\mathbf{c}) & (g \times f) \circ \mathbf{c}_{A,B} = \mathbf{c}_{C,D} \circ (f \times g), \\
(\mathbf{cc}) & \mathbf{c}_{B,A} \circ \mathbf{c}_{A,B} = \mathbf{1}_{A \times B}, \\
(\mathbf{bc} \delta) & (\delta_A \times \mathbf{1}_B) \circ \mathbf{b}_{A,I,B}^\rightarrow = \mathbf{1}_A \times \delta_B \circ \mathbf{c}_{I,B}, \\
(\mathbf{bc6}) & \mathbf{b}_{C,A,B}^\rightarrow \circ \mathbf{c}_{A \times B, C} \circ \mathbf{b}_{A,B,C}^\rightarrow = (\mathbf{c}_{A,C} \times \mathbf{1}_B) \circ \mathbf{b}_{A,C,B}^\rightarrow \circ (\mathbf{1}_A \times \mathbf{c}_{B,C}), \\
(\mathbf{w}) & (f \times f) \circ \mathbf{w}_A = \mathbf{w}_B \circ f, \\
(\delta \mathbf{w}) & \delta_I \circ \mathbf{w}_I = \mathbf{1}_I, \\
(\mathbf{bw}) & \mathbf{b}_{A,A,A}^\rightarrow \circ (\mathbf{1}_A \times \mathbf{w}_A) \circ \mathbf{w}_A = (\mathbf{w}_A \times \mathbf{1}_A) \circ \mathbf{w}_A, \\
(\mathbf{cw}) & \mathbf{c}_{A,A} \circ \mathbf{w}_A = \mathbf{w}_A, \\
(\mathbf{bcw8}) & \mathbf{c}_{A,B,A,B}^m \circ \mathbf{w}_{A \times B} = \mathbf{w}_A \times \mathbf{w}_B, \quad \text{where} \\
& \mathbf{c}_{A,B,C,D}^m \stackrel{\text{df}}{=} \mathbf{b}_{A,C,B \times D}^\rightarrow \circ (\mathbf{1}_A \times (\mathbf{b}_{C,B,D}^\leftarrow \circ (\mathbf{c}_{B,C} \times \mathbf{1}_D) \circ \mathbf{b}_{B,C,D}^\rightarrow)) \circ \mathbf{b}_{A,B,C \times D}^\leftarrow, \\
(\mathbf{k}) & \text{for } f : A \vdash I, f = \mathbf{k}_A, \\
(\delta \mathbf{k} \mathbf{w}) & \delta_A \circ (\mathbf{1}_A \times \mathbf{k}_A) \circ \mathbf{w}_A = \mathbf{1}_A,
\end{array}$$

$$\begin{aligned}
(+1) \quad & \mathbf{1}_A + \mathbf{1}_B = \mathbf{1}_{A+B}, \\
(+2) \quad & (g_1 \circ g_2) + (f_1 \circ f_2) = (g_1 + f_1) \circ (g_2 + f_2), \\
(l^1) \quad & (f_1 + f_2) \circ \mathbf{l}_{A_1, A_2}^1 = \mathbf{l}_{B_1, B_2}^1 \circ f_1, \\
(l^2) \quad & (f_1 + f_2) \circ \mathbf{l}_{A_1, A_2}^2 = \mathbf{l}_{B_1, B_2}^2 \circ f_2, \\
(l) \quad & \text{for } f : \mathbf{O} \vdash A, f = \mathbf{l}_A, \\
(\mathbf{m}) \quad & f \circ \mathbf{m}_A = \mathbf{m}_B \circ (f + f), \\
(\mathbf{lm1}) \quad & \mathbf{m}_A \circ \mathbf{l}_{A, A}^1 = \mathbf{1}_A = \mathbf{m}_A \circ \mathbf{l}_{A, A}^2, \\
(\mathbf{lm2}) \quad & \mathbf{m}_{A+B} \circ (\mathbf{l}_{A, B}^1 + \mathbf{l}_{A, B}^2) = \mathbf{1}_{A+B}, \\
(\rightarrow 1) \quad & \mathbf{1}_A \rightarrow \mathbf{1}_B = \mathbf{1}_{A \rightarrow B}, \\
(\rightarrow 2) \quad & (g_1 \circ g_2) \rightarrow (f_1 \circ f_2) = (g_2 \rightarrow f_1) \circ (g_1 \rightarrow f_1), \\
(\varepsilon 1) \quad & f \circ \varepsilon_{C, A} = \varepsilon_{C, B} \circ (\mathbf{1}_C \times (\mathbf{1}_C \rightarrow f)), \\
(\eta 1) \quad & (\mathbf{1}_C \rightarrow (\mathbf{1}_C \times f)) \circ \eta_{C, A} = \eta_{C, B} \circ f, \\
(\varepsilon 2) \quad & \varepsilon_{B, C} \circ (f \times (\mathbf{1}_B \rightarrow \mathbf{1}_C)) = \varepsilon_{A, C} \circ (\mathbf{1}_A \times (f \rightarrow \mathbf{1}_C)), \\
(\eta 2) \quad & (f \rightarrow (\mathbf{1}_B \times \mathbf{1}_C)) \circ \eta_{B, C} = (\mathbf{1}_A \rightarrow (f \times \mathbf{1}_C)) \circ \eta_{A, C}, \\
(\mathbf{1}\varepsilon\eta) \quad & (\mathbf{1}_A \rightarrow \varepsilon_{A, B}) \circ \eta_{A, A \rightarrow B} = \mathbf{1}_{A \rightarrow B}, \\
(\varepsilon \mathbf{1}\eta) \quad & \varepsilon_{A, A \times B} \circ (\mathbf{1}_A \times \eta_{A, B}) = \mathbf{1}_{A \times B}.
\end{aligned}$$

It is easy to extract the definitions of *symmetric monoidal closed*, *cartesian closed* and *bicartesian* categories from the definition above. This is the first reason to accept the approach above to bicartesian closed categories. Another reason is the sequent system that we are going to deal with, and the process of cut elimination tied to it.

The proof that the above definition is equivalent to the equational definition of bicartesian closed categories given in [9] requires some effort, but we won't go into this matter here.

2 Graphs and g-dinatural transformations

This section is devoted to the notion of g-dinatural transformations. These transformations will serve as morphisms in a functor category whose object are functors of the type $\mathcal{A}_1 \times \mathcal{A}_2 \times \dots \mathcal{A}_n \rightarrow \mathcal{A}$ for some category $\mathcal{A}$, $\mathcal{A}_i \in \{\mathcal{A}, \mathcal{A}^{op}\}$ and arbitrary $n \in \mathbb{N}$. They are always equipped with “graphs” and this explains the letter g in the label g-dinatural. First we define the notion of graph.

For $m \geq 0$ and $n \geq 0$, let M be the set $\{x_1, x_2, \dots, x_m\} + \{y_1, y_2, \dots, y_n\}$ whose elements we call *argument places* (the x 's are *left-hand side* argument places and the y 's are *right-hand side* argument places, and if $m = n = 0$, M is empty). Let G be a finite set and let l be a mapping $l : M \rightarrow \{1, -1\}$, which intuitively denotes the covariance or the contravariance of an argument place. If $l(u) = 1$ we say that u is a *positive argument place* and we write u^+ , and if $l(u) = -1$ we call it *negative argument place* and we write u^- . The elements of the set $V = M \cup G$ are called *vertices*.


Let E be a set of pairs of elements from V that we call *edges*. Let $u \sim v$ means that there is an edge $\{u, v\}$, and let $\simeq$ be the reflexive and transitive closure of $\sim$. Then the equivalence classes of $\simeq$ together with the corresponding edges from E , are called *components*. Let us enumerate these components by $1, 2, \dots, k$, ($k \geq 0$), and let π be the mapping $\pi : V \rightarrow \{1, \dots, k\}$ that maps a vertex from V to the number of its component. We call this function *component classifier*.

For V , l and E as above, the triple (V, l, E) is called *graph* iff the following conditions hold:

1. every vertex belongs to some edge,
2. $\{x_i, x_j\} \in E$ iff $l(x_i) = -l(x_j)$ and x_i, x_j are in the same component,
3. $\{y_i, y_j\} \in E$ iff $l(y_i) = -l(y_j)$ and y_i, y_j are in the same component,

4. $\{x_i, y_j\} \in E$ iff $l(x_i) = l(y_j)$ and x_i, y_j are in the same component,
5. if a component K includes an edge between two argument places, then $K \cap G = \emptyset$; otherwise, $K \cap G = \{g\}$ for some vertex $g \in G$, and for every $u \in K \setminus \{g\}$ the edge $\{u, g\}$ is in $E(K)$.

EXAMPLE 2.1. The following diagram illustrates a graph with 3 components, wher G is a singleton.



For a graph Γ we denote by Γ_i its i -th connectional component. Let Γ_i^+ and Γ_i^- be the sets of positive and negative vertices from Γ_i respectively. Also, for a symbol a we use the abbreviation a^k for the sequence of k occurrences of this symbol.

Let for a single component graph $\Gamma = (V, l, E)$ and a small category $\mathcal{A}$, T and S be functors

$$\begin{aligned} T &: \mathcal{A}^{l(x_1)} \times \dots \times \mathcal{A}^{l(x_m)} \rightarrow \mathcal{A}, \\ S &: \mathcal{A}^{l(y_1)} \times \dots \times \mathcal{A}^{l(y_n)} \rightarrow \mathcal{A}, \end{aligned}$$

where $\mathcal{A}^1 = \mathcal{A}$ and $\mathcal{A}^{-1} = \mathcal{A}^{op}$. Let α be a set

$$\alpha = \{\alpha(A) : T(A^m) \rightarrow S(A^n) \mid A \in Ob(\mathcal{A})\}$$

of morphisms from $\mathcal{A}$ indexed by the set of objects from $\mathcal{A}$. Such a family is called a *transformation*. Then we say that α is a *g-dinatural transformation* from T to S with the graph Γ , which is denoted by $\alpha : T \xrightarrow[\Gamma]{\bullet} S$, if for every pair of objects A, C and every morphism $f : A \rightarrow C$ from $\mathcal{A}$, the following diagram commutes:

$$\begin{array}{ccc} & T\langle C, C \rangle & \xrightarrow{\alpha(C)} S\langle C, C \rangle \\ T\langle f, \mathbf{1}_C \rangle \nearrow & & \searrow S\langle \mathbf{1}_C, f \rangle \\ T\langle A, C \rangle & & S\langle C, A \rangle \\ T\langle \mathbf{1}_A, f \rangle \searrow & & \nearrow S\langle f, \mathbf{1}_A \rangle \\ & T\langle A, A \rangle & \xrightarrow{\alpha(A)} S\langle A, A \rangle \end{array}$$

where $\langle u, v \rangle$ denotes the tuple of arguments with u in positive and v in negative argument places.

Let now Γ be a graph with k ($k > 1$) components, and let

$$\alpha = \{\alpha(A_1, \dots, A_k) : T(A_{\pi(x_1)}, \dots, A_{\pi(x_m)}) \rightarrow S(A_{\pi(y_1)}, \dots, A_{\pi(y_n)}) \mid A_1, \dots, A_k \in Ob(\mathcal{A})\}$$

be a family of morphisms from $\mathcal{A}$ indexed by the set of k -tuples of objects from $\mathcal{A}$. Then we say that α is a *g-dinatural transformation* from T to S with the graph Γ , if for every $k-1$ -tuple $(A_1, \dots, A_{i-1}, A_{i+1}, \dots, A_k)$ of objects from $\mathcal{A}$, the subset


$$\alpha_{A_1, \dots, A_{i-1}, A_{i+1}, \dots, A_k} = \{\alpha(A_1, \dots, A_{i-1}, A, A_{i+1}, \dots, A_k) \mid A \in Ob(\mathcal{A})\}$$

of α is g-dinatural with the graph Γ_i . (All the argument places that are not in Γ_i are parametrized in this case.) This means that a transformation is g-dinatural iff it is g-dinatural in each of its components, or roughly speaking, g-dinaturality is defined componentwise.

EXAMPLE 2.2. Let $\mathcal{C}$ be a cartesian closed category, and let $T : \mathcal{C} \times \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{C}$ and $R : \mathcal{C} \times \mathcal{C}^{op} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ be two functors defined on objects and morphisms of $\mathcal{C}$ by the formulae

$$T(x_1, x_2, x_3) = x_1 \times (x_2 \rightarrow x_3) \quad \text{and} \quad R(z_1, z_2, z_3, z_4) = (z_1 \times (z_2 \rightarrow z_3)) \times z_4.$$

Let Γ be the graph



and let α be the following transformation

$$\{\alpha(A, B) = (\mathbf{1}_{A \times (A \rightarrow B)} \times \varepsilon_{A, B}) \mathbf{w}_{A \times (A \rightarrow B)} : A \times (A \rightarrow B) \vdash (A \times (A \rightarrow B)) \times B \mid A, B \in \text{Ob}(\mathcal{C})\}.$$

Then $\alpha : T \xrightarrow{\bullet \bullet} R$, because for all A, B, C and $f : A \vdash C$ from $\mathcal{C}$ the following two diagrams commute:


$$\begin{array}{ccc} C \times (C \rightarrow B) & \xrightarrow{\alpha(C, B)} & (C \times (C \rightarrow B)) \times B \\ \swarrow f \times (\mathbf{1}_C \rightarrow \mathbf{1}_B) & & \downarrow (\mathbf{1}_C \times (f \rightarrow \mathbf{1}_B)) \times \mathbf{1}_B \\ A \times (C \rightarrow B) & \vdash & (C \rightarrow (A \rightarrow B)) \times B \\ \downarrow \mathbf{1}_A \times (f \rightarrow \mathbf{1}_B) & & \downarrow (f \times (\mathbf{1}_A \rightarrow \mathbf{1}_B)) \times \mathbf{1}_B \\ A \times (A \rightarrow B) & \xrightarrow{\alpha(A, B)} & (A \times (A \rightarrow B)) \times B \end{array}$$

$$\begin{array}{ccc} B \times (B \rightarrow C) & \xrightarrow{\alpha(B, C)} & (B \times (B \rightarrow C)) \times C \\ \swarrow \mathbf{1}_B \times (\mathbf{1}_B \rightarrow f) & & \downarrow (\mathbf{1}_B \times (\mathbf{1}_B \rightarrow \mathbf{1}_C)) \times \mathbf{1}_C \\ B \times (B \rightarrow A) & \vdash & (B \times (B \rightarrow C)) \times C \\ \downarrow \mathbf{1}_B \times (\mathbf{1}_B \rightarrow \mathbf{1}_A) & & \downarrow (\mathbf{1}_B \times (\mathbf{1}_B \rightarrow f)) \times f \\ B \times (B \rightarrow A) & \xrightarrow{\alpha(B, A)} & (B \times (B \rightarrow A)) \times A \end{array}$$

EXAMPLE 2.3. Let β be the following transformation

$$\{\beta(A, B) = ((\mathbf{k}_A \times \mathbf{1}_{A \rightarrow B}) \times \mathbf{1}_B)(\mathbf{1}_{A \times (A \rightarrow B)} \times \varepsilon_{A, B}) \mathbf{w}_{A \times (A \rightarrow B)}(\mathbf{1}_A \times (\mathbf{k}_A \rightarrow \mathbf{1}_B)) \mid A, B \in \mathcal{C}\}$$

between the functors $T : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and $S : \mathcal{C}^{op} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ that are defined by the terms $x_1 \times (\mathbf{I} \rightarrow x_2)$ and $(\mathbf{I} \times (y_1 \rightarrow y_2)) \times y_3$ respectively, for some cartesian closed category $\mathcal{C}$. Then we can show that β is g-dinatural with the graph:



It is obvious how the notion of g-dinaturality extends the traditional notion of dinaturality given in [4]. All that one has to do in order to show that a g-dinatural transformation is already dinatural is to collapse all the argument places of the same sign from a component into one argument place. The main purpose of this extension is to give an answer to the question: “When is the composition of two dinatural transformations dinatural?” The rest of this section is devoted to this problem.

Let $\Phi = (V_\Phi, l_\Phi, E_\Phi)$ where $V_\Phi = \{x_1, \dots, x_m, y_1, \dots, y_n\} \cup G_\Phi$ and $\Psi = (V_\Psi, l_\Psi, E_\Psi)$ where $V_\Psi = \{y_1, \dots, y_n, z_1, \dots, z_p\} \cup G_\Psi$ be two graphs with k_Φ and k_Ψ components, respectively, such that l_Φ and l_Ψ coincide on $\{y_1, \dots, y_n\}$ and that $G_\Phi \cap G_\Psi = \emptyset$. Let

$$T : \mathcal{A}^{l_\Phi(x_1)} \times \mathcal{A}^{l_\Phi(x_2)} \times \dots \times \mathcal{A}^{l_\Phi(x_m)} \rightarrow \mathcal{A}$$

$$S : \mathcal{A}^{l_\Phi(y_1)} \times \mathcal{A}^{l_\Phi(y_2)} \times \dots \times \mathcal{A}^{l_\Psi(y_n)} \rightarrow \mathcal{A}$$

$$R : \mathcal{A}^{l_\Psi(z_1)} \times \mathcal{A}^{l_\Psi(z_2)} \times \dots \times \mathcal{A}^{l_\Psi(z_p)} \rightarrow \mathcal{A}$$

be three functors, and let α and β be two g-dinatural transformations

$$\alpha = \{\alpha(A_1, \dots, A_{k_\Phi}) \mid A_1, \dots, A_{k_\Phi} \in \text{Ob}(\mathcal{A})\} : T \xrightarrow[\Phi]{\bullet\bullet} S$$

$$\beta = \{\beta(B_1, \dots, B_{k_\Psi}) \mid B_1, \dots, B_{k_\Psi} \in \text{Ob}(\mathcal{A})\} : S \xrightarrow[\Psi]{\bullet\bullet} R$$

By the *amalgamation* of Φ and Ψ we mean the couple $(V_\Phi \cup V_\Psi, E_\Phi + E_\Psi)$ denoted by $\Phi + \Psi$. (Note that $\Phi + \Psi$ is not a graph in the sense of the definition above, but we may define its components analogously.)

Let the amalgamation $\Phi + \Psi$ have one component; then we define the graph $\Psi\Phi = (V, l, E)$, i.e. the *composition of the graphs* Φ and Ψ , in the following manner:

- if all x ’s are of the same sign in Φ , which is opposite to the sign of all z ’s in Ψ (this includes the cases when $m = 0$ or $p = 0$), then $V = \{x_1, \dots, x_m\} \cup \{g\}$ and $E = \{\{x_i, g\} \mid 1 \leq i \leq m\} \cup \{\{z_j, g\} \mid 1 \leq j \leq p\}$,
- otherwise, G is empty, $V = \{x_1, \dots, x_m, z_1, \dots, z_p\}$ and $E = \{\{x_i, x_j\} \mid l_\Phi(x_i) = -l_\Phi(x_j)\} \cup \{\{z_i, z_j\} \mid l_\Psi(z_i) = -l_\Psi(z_j)\} \cup \{\{x_i, z_j\} \mid l_\Phi(x_i) = l_\Psi(z_j)\}$.

In both cases, the function l is defined so that its restrictions to $\{x_1, \dots, x_m\}$ and $\{z_1, \dots, z_p\}$ are l_Φ and l_Ψ respectively.

In the case of more than one component in $\Phi + \Psi$, we proceed analogously for each of them to construct a component of the graph $\Psi\Phi$. Since the notion of g-dinaturality is defined componentwise, from now on we consider just the case when $\Phi + \Psi$, and therefore $\Psi\Phi$, has a single component.

Now we define the composition $\beta\alpha$ to be the transformation


$$\{\beta(A^{k_\Psi})\alpha(A^{k_\Phi}) \mid A \in \text{Ob}(\mathcal{A})\}.$$

Our question is: “Is it a g-dinatural transformation with the graph $\Psi\Phi$?”

EXAMPLE 2.4. Let $\mathcal{C}$, T , R be as in Example 2.2 and let $S : \mathcal{A} \times \mathcal{A}^{op} \times \mathcal{A} \times \mathcal{A} \times \mathcal{A}^{op} \times \mathcal{A} \rightarrow \mathcal{A}$ be defined on objects and morphisms by the formula

$$S(y_1, y_2, y_3, y_4, y_5, y_6) = (y_1 \times (y_2 \rightarrow y_3)) \times (y_4 \times (y_5 \rightarrow y_6)).$$

Let Φ be the thin graph and Ψ the thick graph whose amalgamation $\Phi + \Psi$ is given by the diagram



Let β and γ be the transformations


$$\begin{aligned} \{\beta(A, B, C) = \mathbf{w}_{A \times (B \rightarrow C)} \mid A, B, C \in \text{Ob}(\mathcal{C})\} \\ \{\gamma(A, B, C, D, Z) = \mathbf{1}_{A \times (B \rightarrow C)} \times \varepsilon_{D, Z} \mid A, B, C, D, Z \in \text{Ob}(\mathcal{C})\} \end{aligned}$$

Then it is easy to check that $\beta : T \xrightarrow{\ddot{\Phi}} S$, $\gamma : S \xrightarrow{\ddot{\Psi}} R$ and that $\Psi\Phi = \Gamma$ and $\gamma\beta = \alpha$ for Γ and α from Example 2.2.


One may be tempted by these examples to conclude that the composition of g-dinatural transformations is always g-dinatural, as it is the case with natural transformations. This will be proven wrong. However, the category in question may have strong influence on g-dinaturality of the composition of g-dinatural transformations, but we will neglect this possible influence and rely only on the geometry of the underlying graphs. An approach that treats properties intrinsic to a category that are sufficient for dinaturality of a composition of transformations is given in [1].

The next example, although tedious, may serve as a good introduction to what follows.

EXAMPLE 2.5. Let $T : \mathcal{A} \times \mathcal{A}^{op} \times \mathcal{A}^2 \rightarrow \mathcal{A}$, $S : \mathcal{A}^2 \times (\mathcal{A}^{op})^2 \times \mathcal{A} \times (\mathcal{A}^{op})^2 \times \mathcal{A} \rightarrow \mathcal{A}$ and $R : \mathcal{A} \rightarrow \mathcal{A}$ be three functors and $\alpha : T \xrightarrow{\ddot{\Phi}} S$ and $\beta : S \xrightarrow{\ddot{\Psi}} R$ two g-dinatural transformations such that the amalgamation $\Phi + \Psi$ (Ψ is bold) is given by the following diagram



where the components of Φ and Ψ are enumerated by suitable numerals. The composition of Φ and Ψ is given by the diagram




and $\beta\alpha$ is g-dinatural with this graph if the following equation

$$R(\mathbf{1}_C)\beta\alpha(C)T(f, \mathbf{1}_C, f^2) = R(f)\beta\alpha(A)T(\mathbf{1}_A, f, \mathbf{1}_A^2)$$

holds in $\mathcal{A}$ for every A, C and $f : A \rightarrow C$ from this category. We prove this by “travelling” along the amalgamation $\Phi + \Psi$, relying on the definition of $\beta\alpha$, on the functoriality of T, S and R and on the g-dinaturality of α and β . We hope the reader won't be scared with the following a rather long proof in which $(\beta\alpha)$ means reference to the definition of $\beta\alpha$, (T) means reference to functoriality of T , $(\alpha3)$ means reference to g-dinaturality of α in the third component of Φ , etc.

$$\begin{aligned}
& R(\mathbf{1}_C)\beta\alpha(C)T(f, \mathbf{1}_C, f^2) \\
&= R(\mathbf{1}_C)\beta(C^4)\alpha(C^5)T(f, \mathbf{1}_C, f^2) \quad (\beta\alpha) \\
&= R(\mathbf{1}_C)\beta(C^4)\alpha(C^5)T(f, \mathbf{1}_C^3)T(\mathbf{1}_A, \mathbf{1}_C, f^2) \quad (T) \\
&= R(\mathbf{1}_C)\beta(C^4)S(f, \mathbf{1}_C^7)\alpha(A, C^4)T(\mathbf{1}_A, f, \mathbf{1}_C^2)T(\mathbf{1}_A, \mathbf{1}_C, f^2) \quad (\alpha1) \\
&= R(\mathbf{1}_C)\beta(C^3, A)S(\mathbf{1}_A, \mathbf{1}_C^2, f, \mathbf{1}_C^4)\alpha(A, C^4)T(\mathbf{1}_A, f^3) \quad (\beta4), (T) \\
&= R(\mathbf{1}_C)\beta(C^3, A)S(\mathbf{1}_A, \mathbf{1}_C^2, \mathbf{1}_A, f, \mathbf{1}_C^3)\alpha(A, C^2, A, C)T(\mathbf{1}_A, f^3) \quad (\alpha4) \\
&= R(\mathbf{1}_C)\beta(C, A, C, A)S(\mathbf{1}_A, \mathbf{1}_C^2, \mathbf{1}_A^2, f, \mathbf{1}_C^2)\alpha(A, C^2, A, C)T(\mathbf{1}_A, f^3) \quad (\beta2) \\
&= R(\mathbf{1}_C)\beta(C, A, C, A)S(\mathbf{1}_A, \mathbf{1}_C^2, \mathbf{1}_A^2, f, \mathbf{1}_C^2)S(\mathbf{1}_A, \mathbf{1}_C^2, \mathbf{1}_A^2, \mathbf{1}_C^3)\alpha(A, C^2, A, C)T(\mathbf{1}_A^2, f, \mathbf{1}_C) \\
&\quad T(\mathbf{1}_A, f, \mathbf{1}_A, f) \quad (T), (S) \\
&= R(\mathbf{1}_C)\beta(C, A, C, A)S(\mathbf{1}_A, \mathbf{1}_C^2, \mathbf{1}_A^2, f, \mathbf{1}_C^2)S(\mathbf{1}_A, f, \mathbf{1}_C, \mathbf{1}_A^2, \mathbf{1}_C^3)\alpha(A^2, C, A, C) \\
&\quad T(\mathbf{1}_A^3, \mathbf{1}_C)T(\mathbf{1}_A, f, \mathbf{1}_A, f) \quad (\alpha2) \\
&= R(\mathbf{1}_C)\beta(C, A, C, A)S(\mathbf{1}_A, f, \mathbf{1}_C, \mathbf{1}_A^3, \mathbf{1}_C^2)S(\mathbf{1}_A^2, \mathbf{1}_C, \mathbf{1}_A^2, f, \mathbf{1}_C^2)\alpha(A^2, C, A, C) \\
&\quad T(\mathbf{1}_A, f, \mathbf{1}_A, f) \quad (S), (T) \\
&= R(\mathbf{1}_C)\beta(C, A^3)S(\mathbf{1}_A^2, f, \mathbf{1}_A^3, f, \mathbf{1}_C)S(\mathbf{1}_A^2, \mathbf{1}_C, \mathbf{1}_A^2, f, \mathbf{1}_C^2)\alpha(A^2, C, A, C)T(\mathbf{1}_A, f, \mathbf{1}_A, f) \quad (\beta3) \\
&= R(\mathbf{1}_C)\beta(C, A^3)S(\mathbf{1}_A^5, f^2, \mathbf{1}_C)S(\mathbf{1}_A^2, f, \mathbf{1}_A^2, \mathbf{1}_C^3)\alpha(A^2, C, A, C)T(\mathbf{1}_A^3, f)T(\mathbf{1}_A, f, \mathbf{1}_A^2) \quad (T), (S) \\
&= R(\mathbf{1}_C)\beta(C, A^3)S(\mathbf{1}_A^5, f^2, \mathbf{1}_C)S(\mathbf{1}_A^5, \mathbf{1}_C^3)\alpha(A^4, C)T(\mathbf{1}_A^4)T(\mathbf{1}_A, f, \mathbf{1}_A^2) \quad (\alpha3) \\
&= R(\mathbf{1}_C)\beta(C, A^3)S(\mathbf{1}_A^5, f^2, \mathbf{1}_C)\alpha(A^4, C)T(\mathbf{1}_A^4)T(\mathbf{1}_A, f, \mathbf{1}_A^2) \quad (S) \\
&= R(\mathbf{1}_C)\beta(C, A^3)S(\mathbf{1}_A^7, f)\alpha(A^5)T(\mathbf{1}_A, f, \mathbf{1}_A^2) \quad (\alpha5), (T) \\
&= R(f)\beta(A^4)\alpha(A^5)T(\mathbf{1}_A, f, \mathbf{1}_A^2) \quad (\beta1), (S) \\
&= R(f)\beta\alpha(A)T(\mathbf{1}_A, f, \mathbf{1}_A^2) \quad (\beta\alpha)
\end{aligned}$$

This example strengthens the impression that g-dinatural transformations give a g-dinatural transformation in the composition, but could we repeat the above procedure with transformations whose amalgamation of graphs is given below?



Simply, without any further assumptions on the category in question, we can't move along this amalgamation at all.

We shall now examine properties of an amalgamation $\Phi + \Psi$ which guarantee that the composition of $\alpha : T \xrightarrow{\ddot{\Phi}} S$ and $\beta : S \xrightarrow{\ddot{\Psi}} R$ is g-dinatural. For these purposes let Φ and Ψ be as in the definition of amalgamation, and let $\Phi + \Psi$ have one component. We say that $\Phi + \Psi$ *provides g-dinaturality* if for every

category $\mathcal{A}$, for every triple of functors $F : \mathcal{A}^{l_\Phi(x_1)} \times \dots \times \mathcal{A}^{l_\Phi(x_m)} \rightarrow \mathcal{A}$, $G : \mathcal{A}^{l_\Phi(y_1)} \times \dots \times \mathcal{A}^{l_\Phi(y_n)} \rightarrow \mathcal{A}$ and $H : \mathcal{A}^{l_\Psi(z_1)} \times \dots \times \mathcal{A}^{l_\Psi(z_p)} \rightarrow \mathcal{A}$, and for every pair $\gamma : F \xrightarrow[\Phi]{\bullet} G$ and $\delta : G \xrightarrow[\Psi]{\bullet} H$ of g-dinatural transformations, the composition $\delta\gamma$ is g-dinatural from F to H with the graph $\Psi\Phi$. Let $P(\Phi, \Psi)$ denote the property that $\Phi + \Psi$ provides g-dinaturality. To make easier the proof of the main result of this section, we introduce an alternative characterization of $P(\Phi, \Psi)$. In the style of [3] we introduce a free categorial object that will serve as a template for g-dinaturality.

Let $\mathcal{K}_{\Phi, \Psi}$ be the category of structured categories $(\mathcal{A}, F, G, H, \gamma, \delta)$ for $\mathcal{A}, F, G, H, \gamma, \delta$ as above. The morphisms of $\mathcal{K}_{\Phi, \Psi}$ are structure-preserving functors between these categories. The category $\mathcal{K}_{\Phi, \Psi}$ has an equational presentation, as we shall see later; hence, there exists a free object of this category generated by the arrow

$$A \xrightarrow{\mathbf{f}} C.$$

Denote this object by $(\mathcal{D}, T, S, R, \alpha, \beta)$. Its explicit construction will be given soon. The following lemma gives an alternative definition of $P(\Phi, \Psi)$.

LEMMA 2.1. *The amalgamation $\Phi + \Psi$ provides g-dinaturality iff the following diagram*

$$\begin{array}{ccc} & T\langle C, C \rangle & \xrightarrow{\beta(C^{k_\beta})\alpha(C^{k_\alpha})} R\langle C, C \rangle \\ & \nearrow T\langle \mathbf{f}, 1_C \rangle & \searrow R\langle 1_C, \mathbf{f} \rangle \\ T\langle A, C \rangle & & R\langle C, A \rangle \\ & \searrow T\langle 1_A, \mathbf{f} \rangle & \nearrow R\langle \mathbf{f}, 1_A \rangle \\ & T\langle A, A \rangle & \xrightarrow{\beta(A^{k_\beta})\alpha(A^{k_\alpha})} R\langle A, A \rangle \end{array}$$

commutes in $\mathcal{D}$, where $\mathbf{f} : A \rightarrow C$ is the generator of $\mathcal{D}$.

PROOF. The “only if” part of the lemma follows from the definitions of g-dinaturality and of $P(\Phi, \Psi)$. For the “if” part we rely on the universal property of the category $\mathcal{D}$. $\square$

The category $\mathcal{D}$ can be built up from syntactical material in the following manner. The *objects* of $\mathcal{D}$ are freely generated over the set $\{A, C\}$ by the m -ary operation T , the n -ary operation S and the p -ary operation R . We use the schematic letters X, Y and Z , possibly with indices, for elements of $Ob(\mathcal{D})$. The *primitive morphism terms* of $\mathcal{D}$ are

$$\begin{aligned} \mathbf{f} : A &\rightarrow C, & \mathbf{1}_X : X &\rightarrow X, \\ \alpha(Y_1, \dots, Y_{k_\Phi}) &: T(Y_{\pi(x_1)}, \dots, Y_{\pi(x_m)}) &\rightarrow S(Y_{\pi(y_1)}, \dots, Y_{\pi(y_n)}), \\ \beta(Z_1, \dots, Z_{k_\Psi}) &: S(Z_{\pi'(y_1)}, \dots, Z_{\pi'(y_m)}) &\rightarrow R(Z_{\pi'(z_1)}, \dots, Z_{\pi'(z_p)}), \end{aligned}$$

for all objects $X, Y_1, \dots, Y_{k_\Phi}, Z_1, \dots, Z_{k_\Psi}$, where π and π' are component classifiers for Φ and Ψ , respectively.

In the following definitions and equations let F range over the set $\{T, S, R\}$, and let k , depending on F , be the variable for m, n or p respectively.

Morphism terms of $\mathcal{D}$ are defined inductively as follows:

1. primitive morphism terms are morphism terms,
2. if $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ are morphism terms, then $hg : X \rightarrow Z$ is a morphism term,
3. if $\{t_i : X_i \rightarrow Y_i \mid 1 \leq i \leq k \text{ and the } i\text{-th argument place of } F \text{ is positive}\}$ and $\{t_j : Y_j \rightarrow X_j \mid 1 \leq j \leq k, \text{ and the } j\text{-th argument place of } F \text{ is negative}\}$ are two sets of morphism terms, then $F(t_1, \dots, t_m) : F(X_1, \dots, X_m) \rightarrow F(Y_1, \dots, Y_m)$ is a morphism term.

For morphism terms we use the schematic letters g, h, t , possibly primed and with indices, and $\equiv$ is used for identity of terms. *Morphisms* of $\mathcal{D}$ are the equivalence classes of morphism terms modulo congruence generated by the following schematic equations.

Categorical equations

$$(cat1) \quad g\mathbf{1}_X = g = \mathbf{1}_Y g.$$

$$(cat2) \quad t(hg) = (th)g.$$

Functorial equations

$$(F) \quad \text{For morphism terms } g_1, \dots, g_k, h_1, \dots, h_k, t_1, \dots, t_k \text{ such that for every } 1 \leq i \leq k$$

$$t_i \equiv \begin{cases} h_i g_i & \text{; if the } i\text{-th argument place of } F \text{ is positive} \\ g_i h_i & \text{; if the } i\text{-th argument place of } F \text{ is negative} \end{cases},$$

$$F(h_1, \dots, h_k)F(g_1, \dots, g_k) = F(t_1, \dots, t_k).$$

$$(F1) \quad F(\mathbf{1}_{X_1}, \dots, \mathbf{1}_{X_k}) = \mathbf{1}_{F(X_1, \dots, X_k)}.$$

G-dinatural equations

$$(\alpha) \quad \text{For } 1 \leq i \leq k_\Phi \text{ and morphism terms } t : X \rightarrow Y, g_1, \dots, g_m, g'_1, \dots, g'_m, h_1, \dots, h_n, h'_1, \dots, h'_n \text{ such that for } 1 \leq j \leq m$$

$$g_j \equiv \begin{cases} \mathbf{1}_{Z_q} & ; x_j \in \Phi_q \neq \Phi_i \\ t & ; x_j \in \Phi_i^+ \\ \mathbf{1}_Y & ; x_j \in \Phi_i^- \end{cases} \quad g'_j \equiv \begin{cases} \mathbf{1}_{Z_q} & ; x_j \in \Phi_q \neq \Phi_i \\ \mathbf{1}_X & ; x_j \in \Phi_i^+ \\ t & ; x_j \in \Phi_i^- \end{cases}$$

and for $1 \leq j \leq n$

$$h_j \equiv \begin{cases} \mathbf{1}_{Z_q} & ; y_j \in \Phi_q \neq \Phi_i \\ \mathbf{1}_Y & ; y_j \in \Phi_i^+ \\ t & ; y_j \in \Phi_i^- \end{cases} \quad h'_j \equiv \begin{cases} \mathbf{1}_{Z_q} & ; y_j \in \Phi_q \neq \Phi_i \\ t & ; y_j \in \Phi_i^+ \\ \mathbf{1}_X & ; y_j \in \Phi_i^- \end{cases}$$

$$S(h_1, \dots, h_n)\alpha(Z_1, \dots, Z_{i-1}, Y, Z_{i+1}, \dots, Z_{k_\Phi})T(g_1, \dots, g_m) = \\ S(h'_1, \dots, h'_n)\alpha(Z_1, \dots, Z_{i-1}, X, Z_{i+1}, \dots, Z_{k_\Phi})T(g'_1, \dots, g'_m)$$

The equation (β) arises when we replace Φ, m, n, T, α and S in (α) by Ψ, n, p, S, β and R respectively. These three groups of equations are called $\mathcal{K}_{\Phi, \Psi}$ -equations.

The following abbreviations will help us in a syntactical analysis of the category $\mathcal{D}$. Let $[g]$ in a morphism term denote that the morphism term g may occur at that position and let $\mathbb{1}_X$ denote a composition of $q, q \geq 0$, morphism terms $\mathbf{1}_X$. Furthermore we won't use parentheses for composition; hence, from now on equality between morphism terms is taken up to the associativity $(cat2)$.

LEMMA 2.2. *If $g : X \rightarrow Y$ is a morphism term and $X \in \{A, C\}$, then $Y \in \{A, C\}$ and $g \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$. In particular, if $X \equiv C$, then $Y \equiv C$ and $g \equiv \mathbf{1}_C$.*

PROOF. We proceed by induction on the complexity of the morphism term g .

If g is a primitive morphism term, it is neither of the form $\alpha(X_1, \dots, X_{k_\Phi})$ nor $\beta(Y_1, \dots, Y_{k_\Psi})$, since $T(X_{\pi(x_1)}, \dots, X_{\pi(x_m)}) \neq A$, $S(Y_{\pi'(y_1)}, \dots, Y_{\pi'(y_n)}) \neq C$ and $Ob(\mathcal{D})$ is freely generated. Hence, $g \equiv \mathbf{1}_A$ or $g \equiv \mathbf{f}$.

If g is not primitive, then for the same reason as above, g is neither $T(g_1, \dots, g_m)$, nor $S(h_1, \dots, h_n)$, nor $R(t_1, \dots, t_p)$. Hence, g is a composition $g_2 g_1$ for $g_1 : X \rightarrow Z$ and $g_2 : Z \rightarrow Y$. By the inductive hypothesis, since g_1 is of lower complexity than g , $Z \in \{A, C\}$ and $g_1 \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$. Then by the induction hypothesis applied to g_2 , we have $Y \in \{A, C\}$ and $g_2 \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$. Therefore $g \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$, and since $A \neq C$, we claim $g \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$. The second part of the lemma follows from the fact that g is a morphism term. $\square$

Analogously, we can prove:

LEMMA 2.3. *If $g : X \rightarrow Y$ is a morphism term and $Y \in \{A, C\}$, then $X \in \{A, C\}$ and $g \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$.*

Let $\mathbf{T}$ abbreviate a composition of q , $q \geq 0$, morphism terms of the form $\mathbb{1}T(\mathbb{1}_C[\mathbf{f}]\mathbb{1}_A, \dots, \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A)\mathbb{1}$, and let $\mathbf{S}$ and $\mathbf{R}$ mean the same for S and R instead of T respectively. Denote by $\mathcal{M}$ the set of morphism terms of the form

$$\mathbf{R}\beta(Y_1, \dots, Y_{k_\Psi})\mathbf{S}\alpha(X_1, \dots, X_{k_\Phi})\mathbf{T}$$

for $X_1, \dots, X_{k_\Phi}, Y_1, \dots, Y_{k_\Psi} \in \{A, C\}$, whose type is $T\langle A, C \rangle \rightarrow R\langle C, A \rangle$.

LEMMA 2.4. *The set $\mathcal{M}$ is closed under equality.*

PROOF. A substitution of equalities according to the categorial and functorial equations doesn't change the form of a term from $\mathcal{M}$. Substitutions of equalities according to the "limit" cases of (α) and (β) cause suspicion. Such is, for example, the case of substitution according to (α) when $\Phi_i^+ \cap \{x_1, \dots, x_m\} = \Phi_i^- \cap \{y_1, \dots, y_n\} = \emptyset$. If g' is a term obtained by such a substitution from an $\mathcal{M}$ morphism term g , then an arbitrary morphism term $t : X \rightarrow C$ may occur as an argument of T and S , and this X may occur as an argument of α in g' . However, Lemma 2.3 guarantees that then $X \in \{A, C\}$ and $t \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$, hence g' remains in $\mathcal{M}$. We deal with the other limit cases analogously, referring to Lemmata 2.2 and 2.3 when necessary. Nonlimit cases of substitution according to (α) and (β) are obviously harmless. $\square$

EXAMPLE 2.6. Let Φ and Ψ be as in Example 2.4. Consider the morphism term

$$h \equiv S(\mathbf{1}_A^5, \mathbf{1}_C^3)\alpha(A^4, C)T(\mathbf{1}_A^4).$$

By g-dinaturality of α in the third component of Φ , it is equal to

$$h' = S(\mathbf{1}_A^2, t, \mathbf{1}_A^2, \mathbf{1}_C^3)\alpha(A^2, X, A, C)T(\mathbf{1}_A^3, t)$$

for some $t : A \rightarrow X$. Then by Lemma 2.2, $X \in \{A, C\}$ and $t \equiv \mathbb{1}_C[\mathbf{f}]\mathbb{1}_A$ which is enough for a term to remain in $\mathcal{M}$ after the substitution of h' for h in it.

Let (nat) denote the equation

$$R\langle \mathbf{1}_C, \mathbf{f} \rangle \beta(C^{k_\Psi}) \alpha(C^{k_\Phi}) T\langle \mathbf{f}, \mathbf{1}_C \rangle = R\langle \mathbf{f}, \mathbf{1}_A \rangle \beta(A^{k_\Psi}) \alpha(A^{k_\Phi}) T\langle \mathbf{1}_A, \mathbf{f} \rangle.$$

It is clear that (nat) means commutativity of the diagram from Lemma 2.1, and therefore

$$(nat) \Leftrightarrow P(\Phi, \Psi).$$

So to prove that $P(\Phi, \Psi)$ is decidable we may use a normalization procedure in a rewrite system corresponding to the equational theory of $\mathcal{M}$. Actually, we have two notions of reductions. The first one is called *CF* (*categorial-functorial reduction*), and its redexes and contracta are the following

<i>CF</i> step	<i>redex</i>	<i>contractum</i>
(1)	$g\mathbf{1}$	g
(2)	$\mathbf{1}g$	g
(3)	$F(h_1, \dots, h_k)F(g_1, \dots, g_k)$	$F(t_1, \dots, t_k)$
(4)	$F(\mathbf{1}_{X_1}, \dots, \mathbf{1}_{X_k})$	$\mathbf{1}_{F(D_1, \dots, D_k)}$

In the last two steps F , k , g 's, h 's and t 's satisfy the conditions from the functorial equations above.


Since a CF redex and the corresponding contractum are equal, by Lemma 2.4 we have that a term remains in $\mathcal{M}$ after a CF reduction.

By the following lemma we have that each morphism term g from $\mathcal{M}$ has a unique CF -normal form, which we denote by $CF(g)$.

LEMMA 2.5. *CF is strongly normalizing and weakly Church-Rosser.*

PROOF. For strong normalization it is enough to note that a CF contractum is of lower complexity than the corresponding redex.

The only interesting cases in proving that CF is weakly Church-Rosser are the following (the other cases of ramification, roughly speaking, commute):



and the analogous case starting with $F(\mathbf{1}^k)F(g_1, \dots, g_k)$.

Let $\mathcal{M}_0$ be the set of morphism terms from $\mathcal{M}$ in CF normal form. Henceforth we use the abbreviations $\vec{X}, \vec{Y}, \vec{Z}, \dots$ for tuples of elements from the set $\{A, C\}$ and $\vec{g}, \vec{h}, \vec{t}, \dots$ for tuples of elements from the set $\{\mathbf{1}_A, \mathbf{1}_C, \mathbf{f}\}$. From now on, a subterm in square brackets occurs only if at least one of its arguments is $\mathbf{f}$. With this notation, we have that each member of $\mathcal{M}_0$ is of the shape

$$[R(\vec{t})]\beta(\vec{Y})[S(\vec{h})]\alpha(\vec{X})[T(\vec{g})].$$

The second notion of reduction, called *D-reduction*, where D stands for dinatural, is defined on morphism terms from $\mathcal{M}_0$. A peculiarity of this reduction is that it is applicable only to the entire term as the redex, and not to its subterms. Otherwise, it would be possible to get out of $\mathcal{M}_0$.

For every i , $1 \leq i \leq k_\Phi$ and $\vec{X}, \vec{Y}, \vec{g}, \vec{h}, \vec{t}$ such that both sets $\{h_j \mid y_j \in \Phi_i^-\}$ and $\{g_j \mid x_j \in \Phi_i^+\}$ are subsets of the singleton $\{\mathbf{f}\}$, the morphism term from $\mathcal{M}_0$ of the following form (whose type must be $T\langle A, C \rangle \rightarrow R\langle C, A \rangle$)

$$[R(\vec{t})]\beta(\vec{Y})[S(\vec{h})]\alpha(X_1, \dots, X_{i-1}, C, X_{i+1}, \dots, X_{k_\Phi})[T(\vec{g})]$$

is a redex and

$$[R(\vec{t})]\beta(\vec{Y})[S(\vec{h}')] \alpha(X_1, \dots, X_{i-1}, A, X_{i+1}, \dots, X_{k_\Phi})[T(\vec{g}')]]$$

$$\text{where } g'_j \equiv \begin{cases} g_j & ; x_j \notin \Phi_i \\ \mathbf{1}_A & ; x_j \in \Phi_i^+ \\ \mathbf{f} & ; x_j \in \Phi_i^- \end{cases} \quad \text{and} \quad h'_j \equiv \begin{cases} h_j & ; y_j \notin \Phi_i \\ \mathbf{f} & ; y_j \in \Phi_i^+ \\ \mathbf{1}_A & ; y_j \in \Phi_i^- \end{cases}$$

is the contractum of an (α_i) -step of D reduction. Note that both the redex and the contractum of this step are in $\mathcal{M}_0$. It follows from this fact, together with Lemmata 2.2 and 2.3, that $\{h_j \mid y_j \in \Phi_i^+\}$ and $\{g_j \mid x_j \in \Phi_i^-\}$ are subsets of $\{\mathbf{1}_C\}$.

Analogously, for a fixed $1 \leq i \leq k_\Psi$, we introduce (β_i) -steps of D reduction whose redexes are terms from $\mathcal{M}_0$ of the form

$$[R(\vec{t})]\beta(Y_1, \dots, Y_{i-1}, C, Y_{i+1}, \dots, Y_{k_\Psi})[S(\vec{h})]\alpha(\vec{X})[T(\vec{g})]$$

with both sets $\{t_j \mid z_j \in \Psi_i^-\}$ and $\{h_j \mid y_j \in \Psi_i^+\}$ being subsets of the singleton $\{\mathbf{f}\}$; the corresponding contractum is the morphism term

$$[R(\vec{t'})]\beta(Y_1, \dots, Y_{i-1}, A, Y_{i+1}, \dots, Y_{k_\Psi})[S(\vec{h'})]\alpha(\vec{X})[T(\vec{g})],$$

$$\text{where } h'_j \equiv \begin{cases} h_j & ; y_j \notin \Psi_i \\ \mathbf{1}_A & ; y_j \in \Psi_i^+ \\ \mathbf{f} & ; x_j \in \Psi_i^- \end{cases} \quad \text{and} \quad t'_j \equiv \begin{cases} t_j & ; z_j \notin \Psi_i \\ \mathbf{f} & ; z_j \in \Psi_i^+ \\ \mathbf{1}_A & ; z_j \in \Psi_i^- \end{cases}$$

EXAMPLE 2.7. For Φ and Ψ as in Example 2.4 we have the following (α_1) step of D reduction.

$$R(\mathbf{1}_C)\beta(C^4)\alpha(C^5)T(\mathbf{f}, \mathbf{1}_C, \mathbf{f}^2) \rightsquigarrow R(\mathbf{1}_C)\beta(C^4)S(\mathbf{f}, \mathbf{1}_C^7)\alpha(A, C^4)T(\mathbf{1}_A, \mathbf{f}^3).$$

By the following lemma we establish the uniqueness of D normal form of a morphism term from $\mathcal{M}_0$. We denote the D normal form of g by $D(g)$.

LEMMA 2.6. *D is strongly normalizing and weakly Church-Rosser.*

PROOF. The strong normalization property follows from the fact that every reduction step decreases the number of C 's as arguments of α and β . For the proof that D is weakly Church-Rosser, we rely on the following facts:

- reduction steps (α_i) and (α_j) ((β_i) and (β_j)) commute for $i \neq j$, since connectional components of a graph are disjoint,
- if a term from $\mathcal{M}_0$ is the redex of (α_i) and (β_j) reduction steps, then there is no q , $1 \leq q \leq n$, for which y_q is in both Φ_i and Ψ_j . This is because from the initial assumption it follows that $y_q \in \Phi_i^+$ claims $h_q \equiv \mathbf{1}_C$ and $y_q \in \Psi_j^+$ claims $h_q \equiv \mathbf{f}$ and from the similar reason y_q can't be a negative vertex in $\Phi_i \cap \Psi_j$. Hence, the reduction steps (α_i) and (β_j) act on disjoint sets of arguments of T , S , R , α and β and therefore commute. $\square$

We shall find Lemmata 2.5 and 2.6 very useful for

THEOREM 2.1. *Equality in $\mathcal{M}$ is decidable.*

PROOF. It is enough to show that for two morphism terms g_1 and g_2 from $\mathcal{M}$ the following equivalence holds:

$$g_1 = g_2 \quad \text{iff} \quad D(CF(g_1)) \equiv D(CF(g_2))$$

The *if* part of this equivalence is trivial since all the reductions are covered by our equations (CF reductions are covered by categorial and functorial equations and for D reductions we need all $\mathcal{K}_{\Phi, \Psi}$ equations).

To prove the *only if* part, we rely on the equality axioms (reflexivity, symmetry, transitivity and congruence), and we assume that g_2 is the result of a substitution of a term for a subterm of g_1 according to a $\mathcal{K}_{\Phi, \Psi}$ equation. (By the equality axioms, we must have a chain of morphism terms $g_1 \equiv h_0 = h_1 = \dots = h_q \equiv g_2$ such that for adjacent terms, one is obtained from the other by a substitution described above.) If the equation in question is a categorial or functorial equation, then by Lemma 2.5, we have that $CF(g_1) \equiv CF(g_2)$; hence $D(CF(g_1)) \equiv D(CF(g_2))$. If we deal with a dinatural equation, then it is clear that we need just one step of D reduction to reduce $CF(g_1)$ to $CF(g_2)$ or vice versa, and therefore, by Lemma 2.6, $D(CF(g_1)) \equiv D(CF(g_2))$. $\square$

COROLLARY *The property $P(\Phi, \Psi)$ is decidable.*

Let us transform the equation (nat) by deleting superfluous subterms, if necessary, to obtain the following equation

$$(cfnat) \quad [R(\mathbf{1}_C, \mathbf{f})]\beta(C^{k_\Psi})\alpha(C^{k_\Phi})[T(\mathbf{f}, \mathbf{1}_C)] = [R(\mathbf{f}, \mathbf{1}_A)]\beta(A^{k_\Psi})\alpha(A^{k_\Phi})[T(\mathbf{1}_A, \mathbf{f})].$$

It is easy to see that the left-hand side (*LHS*) and the right-hand side (*RHS*) of (*cfnat*) are in *CF* normal form. Moreover, *RHS* is in *D* normal form too. Therefore, the property $P(\Phi, \Psi)$ is equivalent to

$$D(LHS) \equiv RHS.$$

We use this equivalence in order to establish some geometrical conditions of the amalgamation $\Phi + \Psi$, which are equivalent to $P(\Phi, \Psi)$. For this reason we introduce the following auxiliary notation. For a graph Γ and $v \in V_\Gamma \setminus G_\Gamma$, let Γ_v be the set $\{w \in \Gamma_{\pi(v)} \setminus G_\Gamma \mid \{v, w\} \notin E_\Gamma\}$, and let Γ'_v be the set $\{w \in \Gamma_{\pi(v)} \setminus G_\Gamma \mid \{v, w\} \in E_\Gamma\}$. With this notation, in Example 2.4, we have $\Psi_{y_2} = \{y_2\}$, $\Psi'_{y_2} = \{y_3, y_7\}$, $\Phi_{x_1} = \{x_1\}$, $\Phi'_{x_1} = \{x_2, y_1\}$, $\Phi_{x_4} = \{x_4, y_3\}$, $\Phi'_{x_4} = \emptyset$, etc.

LEMMA 2.7. *For a positive y_i let a morphism term from $\mathcal{M}_0$ in which the i -th argument of S is $\mathbf{1}_C$, reduce by a sequence of D reductions to a term in which this argument is $\mathbf{f}$. Then this sequence of reductions includes a step in whose redex all the argument places from Φ'_{y_i} are occupied by $\mathbf{f}$ and the i -th argument of S is $\mathbf{1}_C$.*

PROOF. Suppose that

$$[R(\vec{t}^0)]\beta(Y^0)[S(\vec{h}^0)]\alpha(\vec{X}^0)[T(\vec{g}^0)] \rightsquigarrow [R(\vec{t}^1)]\beta(Y^1)[S(\vec{h}^1)]\alpha(\vec{X}^1)[T(\vec{g}^1)] \rightsquigarrow \dots \rightsquigarrow [R(\vec{t}^q)]\beta(Y^q)[S(\vec{h}^q)]\alpha(\vec{X}^q)[T(\vec{g}^q)],$$

is the shortest sequence of D reductions for which the lemma fails. Hence, $h_i^0 \equiv \mathbf{1}_C$ and $h_i^q \equiv \mathbf{f}$. We claim that $h_i^1 \not\equiv \mathbf{1}_C$, otherwise we would have a shorter sequence than the initial for which the lemma fails. Also, h_i^1 is not $\mathbf{f}$; otherwise, the first reduction step requires all the argument places from Φ'_{y_i} in the redex to be occupied by $\mathbf{f}$, which together with $h_i^0 \equiv \mathbf{1}_C$ contradicts the assumption that the lemma fails. Eventually, $h_i^1 \equiv \mathbf{1}_A$ is impossible because there is no D reduction step transforming $\mathbf{1}_C$ to $\mathbf{1}_A$ directly. Hence, the lemma holds, since we have exhausted all the possibilities for h_i^1 . $\square$

LEMMA 2.8. *For a positive y_i let a morphism term from $\mathcal{M}_0$ in which the i -th argument of S belongs to the set $\{\mathbf{f}, \mathbf{1}_C\}$, reduce by a sequence of D reductions to a term in which this argument is $\mathbf{1}_A$. Then this sequence of reductions includes a step in whose redex all the argument places from Ψ_{y_i} are occupied by $\mathbf{f}$ and in whose contractum all the argument places from Ψ_{y_i} are occupied by $\mathbf{1}_A$ and all the argument places from Ψ'_{y_i} are occupied by $\mathbf{f}$.*

PROOF. Let again

$$[R(\vec{t}^0)]\beta(Y^0)[S(\vec{h}^0)]\alpha(\vec{X}^0)[T(\vec{g}^0)] \rightsquigarrow [R(\vec{t}^1)]\beta(Y^1)[S(\vec{h}^1)]\alpha(\vec{X}^1)[T(\vec{g}^1)] \rightsquigarrow \dots \rightsquigarrow [R(\vec{t}^q)]\beta(Y^q)[S(\vec{h}^q)]\alpha(\vec{X}^q)[T(\vec{g}^q)],$$

be a shortest sequence of reductions for which the lemma fails. Note that $h_i^0 \in \{\mathbf{f}, \mathbf{1}_C\}$ and $h_i^q \equiv \mathbf{1}_A$. Now h_i^1 is neither $\mathbf{f}$ nor $\mathbf{1}_C$; otherwise we would have a shorter sequence for which the lemma fails. Also, $h_i^1 \not\equiv \mathbf{1}_A$; otherwise, the first reduction step requires arguments in the redex and in the contractum such that it contradicts the assumption that the lemma fails. $\square$

We can prove the following two lemmata analogously.

LEMMA 2.9. *For a negative y_i let a morphism term from $\mathcal{M}_0$ in which the i -th argument of S is $\mathbf{1}_C$, reduce by a sequence of D reductions to a term in which this argument is $\mathbf{f}$. Then this sequence of reductions includes a step in whose redex all the argument places from Ψ'_{y_i} are occupied by $\mathbf{f}$ and the i -th argument of S is $\mathbf{1}_C$.*

LEMMA 2.10. *For a negative y_i let a morphism term from $\mathcal{M}_0$ in which the i -th argument of S belongs to the set $\{\mathbf{f}, \mathbf{1}_C\}$, reduce by a sequence of D reductions to a term in which this argument is $\mathbf{1}_A$. Then this sequence of reductions includes a step in whose redex all the argument places from Φ_{y_i} are occupied by $\mathbf{f}$ and in whose contractum all the argument places from Φ_{y_i} are occupied by $\mathbf{1}_A$ and all the argument places from Φ'_{y_i} are occupied by $\mathbf{f}$.*

In the sequel we also refer to the propositions concerning an x or a z vertex instead of y_i , which are analogous to the last four lemmata.

We are ready to define a geometrical criterion for $P(\Phi, \Psi)$. Let $v_1, v_2, \dots, v_q$ be a sequence of vertices and let $e_1, e_2, \dots, e_{q-1}$ be a sequence of edges from $\Phi + \Psi$ such that $e_1 = \{v_1, v_2\}$, $e_2 = \{v_2, v_3\}$, etc., and such that for each pair of adjacent edges, one belongs to E_Φ and the other to E_Ψ . We call such a pair of sequences an *alternating chain*. If $v_1 = v_q$, then the alternating chain is called an *alternating loop*. Note that in the latter case, the edges e_1 and e_{q-1} are not in the same graph, and the name alternating loop is still justified. Here is a necessary condition for $P(\Phi, \Psi)$.

LEMMA 2.11. *If $\Phi + \Psi$ provides g -dinaturality, then there are no alternating loops in it.*

PROOF. From the definition of graph it follows that the sequence of vertices in an alternating loop consists of an even number of mutually distinct y 's. Suppose now that $P(\Phi, \Psi)$ holds but that $\Phi + \Psi$ includes an alternating loop. For the sake of clarity we use the simplest case with the loop whose vertices are y_i^+ and y_j^- and whose edges are $e_1 = \{y_i, y_j\} \in E_\Phi$ and $e_2 = \{y_i, y_j\} \in E_\Psi$. In all the other cases we can proceed analogously.

By the corollary of Theorem 2.1 and by the assumption $P(\Phi, \Psi)$ we have that the term

$$g_1 \equiv [R\langle \mathbf{1}_C, \mathbf{f} \rangle] \beta(C^{k_\Psi}) \alpha(C^{k_\Phi}) [T\langle \mathbf{f}, \mathbf{1}_C \rangle]$$

reduces by a D reduction to the term

$$g_2 \equiv [R\langle \mathbf{f}, \mathbf{1}_A \rangle] \beta(A^{k_\Psi}) \alpha(A^{k_\Phi}) [T\langle \mathbf{1}_A, \mathbf{f} \rangle].$$

By Lemma 2.8, this reduction must be of the form

$$g_1 \rightsquigarrow \dots \rightsquigarrow g_3 \rightsquigarrow \dots \rightsquigarrow g_2$$

with g_3 an $\mathcal{M}_0$ morphism term whose i -th argument of S is $\mathbf{f}$. Then by Lemma 2.7 this reduction must be of the form

$$g_1 \rightsquigarrow \dots \rightsquigarrow g_4 \rightsquigarrow \dots \rightsquigarrow g_3 \rightsquigarrow \dots \rightsquigarrow g_2$$

with g_4 an $\mathcal{M}_0$ morphism term whose j -th argument place of S is $\mathbf{f}$. By Lemma 2.9, the reduction must be of the form

$$g_1 \rightsquigarrow \dots \rightsquigarrow g_5 \rightsquigarrow \dots \rightsquigarrow g_4 \rightsquigarrow \dots \rightsquigarrow g_3 \rightsquigarrow \dots \rightsquigarrow g_2$$

with the i -th argument of S being $\mathbf{f}$ in g_5 . Now we can repeat this procedure endlessly which contradicts to the finiteness of the reduction $\square$

The necessity of our geometrical condition for $P(\Phi, \Psi)$ is of rather smaller practical interest for the purpose of proving dinaturality of transformations. It can be used in a construction of a countermodel for the dinaturality of composition. However, the other direction of the lemma above is much more useful and we are going to prove it now. For this purposes we define the following binary relation $<_\Gamma$ in the set of the argument places of a graph Γ : every positive left-hand side argument place u is in the relation $<_\Gamma$ with every element of Γ'_u and every negative right-hand side argument place v is in the relation $<_\Gamma$ with every member of Γ'_v . For an amalgamation $\Phi + \Psi$ let $<$ be the union of $<_\Phi$ and $<_\Psi$. By this definition, we have the following chains arranged by $<$ in Example 2.4.

$$\begin{array}{c} x_1 < x_2 \\ x_1 < y_1 < y_4 < y_5 < y_6 < y_8 < z_1 \\ x_3 < y_2 < y_7 < y_8 < z_1 \\ x_3 < y_2 < y_3 \\ x_4 \end{array}$$

LEMMA 2.12. *If there are no alternating loops in $\Phi + \Psi$, then this amalgamation provides g-dinaturality.*

PROOF. Let $\leq$ be the reflexive and transitive closure of $<$ defined as above in the set of argument places from $\Phi \cup \Psi$. This set is partially ordered by $\leq$ because of the absence of alternating loops in $\Phi + \Psi$. Suppose now that $P(\Phi + \Psi)$ fails; i.e., for the equality (*cfnat*) we have

$$D(LHS) \equiv [R(\vec{t})]\beta(\vec{Y})[S(\vec{h})]\alpha(\vec{X})[T(\vec{g})] \neq RHS.$$

Hence, at least one of the following cases must occur in $D(LHS)$.

1. An argument of R , S or T is $\mathbf{1}_C$.
2. For some i such that x_i is positive, g_i is $\mathbf{f}$.
3. For some i such that x_i is negative, g_i is $\mathbf{1}_A$.
4. For some i , h_i is $\mathbf{f}$.
5. For some i such that z_i is positive, t_i is $\mathbf{1}_A$.
6. For some i such that z_i is negative, t_i is $\mathbf{f}$.

Cases 3. and 5. are impossible since the reduction preserves types of morphism terms.

Suppose now that we have Case 1. In the ordering $\leq$, let an argument place v be minimal such that it is occupied by $\mathbf{1}_C$ in $D(LHS)$. The vertex v is neither of the form x^+ nor z^- for the same reason as above. Suppose that $v \equiv x_i^-$. We deal with the other cases analogously. The set Φ'_{x_i} couldn't be empty; otherwise, $D(LHS)$ is the redex of an (α_i) step of D reduction. An argument place from Φ'_{x_i} couldn't be occupied by $\mathbf{1}_C$ in $D(LHS)$, since for every $v \in \Phi'_{x_i}$, $v < x_i$. If all the argument places from Φ'_{x_i} are occupied by $\mathbf{f}$ in $D(LHS)$, then it is not in normal form. If an argument place from Φ'_{x_i} is occupied by $\mathbf{1}_A$ in $D(LHS)$, then by an analogue of Lemma 2.8 (concerning the vertex x_i instead of y_i) the reduction

$$LHS \rightsquigarrow \dots \rightsquigarrow D(LHS)$$

includes a step in whose redex the i -th argument of T is $\mathbf{f}$. Since there is no reduction transforming $\mathbf{f}$ into $\mathbf{1}_C$, and since $g_i \equiv \mathbf{1}_C$ in $D(LHS)$, this is impossible. Therefore, Case 1 leads to a contradiction.

Suppose now we have Case 2. As we have just seen, Case 1. doesn't obtain. If all the argument places from Φ_{x_i} are occupied by $\mathbf{f}$ in $D(LHS)$, then it is not a D normal form. Let $x_j \in \Phi_{x_i}$ be occupied by $\mathbf{1}_A$. The other cases are dealt with analogously. By an analogue of Lemma 2.8 (concerning x_j instead of y_i) the reduction

$$LHS \rightsquigarrow \dots \rightsquigarrow D(LHS)$$

includes a step in whose contractum all the argument places from Φ_{x_i} are occupied by $\mathbf{1}_A$. Hence x_i is occupied by $\mathbf{1}_A$ in this morphism term. Since no reduction transforms $\mathbf{1}_A$ into $\mathbf{f}$, this is impossible. With cases 4. and 6. we deal analogously. $\square$

Composing the previous two lemmata, we obtain the main result of the section.

THEOREM 2.2. $P(\Phi, \Psi) \Leftrightarrow \Phi + \Psi$ *doesn't include alternating loops.*

Note that this theorem considers just a single component amalgamation $\Phi + \Psi$, but as it was mentioned earlier, this result holds universally since the notion of g-dinaturality is defined component-wise. Also, we have reduced our considerations to functors with arguments from one category. The generalization is trivial but it would complicate the notation which is already, by our opinion, at the limit of acceptability.

It is time now to compare this result with a classical one from [5], which has served as an inspiration for our Theorem 2.2. However, the basis of [5] (definitions of graph and naturality) was created to fit applications involving symmetric monoidal closed categories (cf. [8]), and it is obvious that we have

here in mind a more involved case of bicartesian closed categories. We believe that our result may be applicable beyond this limitation. It is easy to see how the part of our theorem concerning sufficiency of the given condition for $P(\Phi, \Psi)$ captures the main result given in [5]. The lack of closed curves in $\Phi + \Psi$, which was taken there as sufficient for $P(\Phi, \Psi)$, has as a trivial consequence the lack of alternating loops. In fact these two conditions are equivalent in the scope of the restricted definition of graph given in [5], since there are no points of ramification in $\Phi + \Psi$. However, in our context the presence of closed curves is harmless for dinaturality by itself; we must instead rely on the absence of *alternating* loops in amalgamations in order to guarantee dinaturality.

3 Bicartesian closed canonical transformations

By a *bicartesian closed canonical* (also called *allowable*) *transformation* in a bicartesian closed category $\mathcal{B}$ we mean a set of morphisms from this category indexed by the objects from $\mathcal{B}$, defined in terms of the special morphisms and the morphism operations from the definition given in Section 1. Formally, it can be defined in the following manner.

Let $\mathcal{F}_{\mathcal{B}}$ be the category whose objects are functors of types $\mathcal{B}^0 \rightarrow \mathcal{B}$, where $\mathcal{B}^0$ is the trivial category $1_* : * \rightarrow *$, or $\mathcal{B}^{l_1} \times \dots \times \mathcal{B}^{l_m} \rightarrow \mathcal{B}$ for $m \geq 0$ and $l_i \in \{-1, 1\}$. We define $Ob(\mathcal{F}_{\mathcal{B}})$ inductively by

$$\begin{aligned} 1_{\mathcal{B}} : \mathcal{B} &\rightarrow \mathcal{B} \quad \in Ob(\mathcal{F}_{\mathcal{B}}), \\ I : \mathcal{B}^0 &\rightarrow \mathcal{B} \quad (I(*) = I) \quad \in Ob(\mathcal{F}_{\mathcal{B}}), \\ O : \mathcal{B}^0 &\rightarrow \mathcal{B} \quad (O(*) = O) \quad \in Ob(\mathcal{F}_{\mathcal{B}}). \end{aligned}$$

If $F : \mathcal{B}^{l_1} \times \dots \times \mathcal{B}^{l_m} \rightarrow \mathcal{B}$ and $G : \mathcal{B}^{l_{m+1}} \times \dots \times \mathcal{B}^{l_{m+n}} \rightarrow \mathcal{B}$ are in $Ob(\mathcal{F}_{\mathcal{B}})$, then $F \otimes G : \mathcal{B}^{l_1} \times \dots \times \mathcal{B}^{l_m} \times \mathcal{B}^{l_{m+1}} \times \dots \times \mathcal{B}^{l_{m+n}} \rightarrow \mathcal{B}$ ($F \otimes G(x_1, \dots, x_{m+n}) = F(x_1, \dots, x_m) \times G(x_{m+1}, \dots, x_{m+n})$), $F \oplus G : \mathcal{B}^{l_1} \times \dots \times \mathcal{B}^{l_m} \times \mathcal{B}^{l_{m+1}} \times \dots \times \mathcal{B}^{l_{m+n}} \rightarrow \mathcal{B}$ ($F \oplus G(x_1, \dots, x_{m+n}) = F(x_1, \dots, x_m) + G(x_{m+1}, \dots, x_{m+n})$) and $F \rightarrow G : \mathcal{B}^{-l_1} \times \dots \times \mathcal{B}^{-l_m} \times \mathcal{B}^{l_{m+1}} \times \dots \times \mathcal{B}^{l_{m+n}} \rightarrow \mathcal{B}$ ($F \rightarrow G(x_1, \dots, x_{m+n}) = F(x_1, \dots, x_m) \rightarrow G(x_{m+1}, \dots, x_{m+n})$) are in $Ob(\mathcal{F}_{\mathcal{B}})$.

The set of *canonical transformations* that we define below will be the set of morphisms from $\mathcal{F}_{\mathcal{B}}$. Each canonical transformation is a set of $\mathcal{B}$ morphisms indexed by tuples of objects from $\mathcal{B}$, together with a graph defined as in Section 2. First we define *primitive canonical transformations* for every $F : \mathcal{B}^{l_1} \times \dots \times \mathcal{B}^{l_m} \rightarrow \mathcal{B}$, $G : \mathcal{B}^{l_{m+1}} \times \dots \times \mathcal{B}^{l_{m+n}} \rightarrow \mathcal{B}$ and $H : \mathcal{B}^{l_{m+n+1}} \times \dots \times \mathcal{B}^{l_{m+n+p}} \rightarrow \mathcal{B}$ from $Ob(\mathcal{F}_{\mathcal{B}})$.

$\mathbf{1}_F = \{\mathbf{1}_{F(\vec{A})} \mid \vec{A} \in (Ob(\mathcal{B}))^m\}$ is a primitive canonical transformation from F to F whose graph consists of vertices $x_1^{l_1}, \dots, x_m^{l_m}, y_1^{l_1}, \dots, y_m^{l_m}$ and edges $\{x_1, y_1\}, \dots, \{x_m, y_m\}$.

$\delta_F = \{\delta_{F(\vec{A})} \mid \vec{A} \in (Ob(\mathcal{B}))^m\}$ is a primitive canonical transformation from $F \otimes I$ to F whose graph is identical to the graph of $\mathbf{1}_F$.

$\mathbf{c}_{F,G} = \{\mathbf{c}_{F(\vec{A}), G(\vec{B})} \mid \vec{A} \in (Ob(\mathcal{B}))^m, \vec{B} \in (Ob(\mathcal{B}))^n\}$ is a primitive canonical transformation from $F \otimes G$ to $G \otimes F$ with the graph whose vertices are $x_1^{l_1}, \dots, x_m^{l_m}, x_{m+1}^{l_{m+1}}, \dots, x_{m+n}^{l_{m+n}}, y_1^{l_{m+1}}, \dots, y_n^{l_{m+n}}, y_{n+1}^{l_1}, \dots, y_{n+m}^{l_m}$ and whose edges are $\{x_1, y_{n+1}\}, \dots, \{x_m, y_{n+m}\}, \{x_{m+1}, y_1\}, \dots, \{x_{m+n}, y_n\}$.

$\mathbf{w}_F = \{\mathbf{w}_{F(\vec{A})} \mid \vec{A} \in (Ob(\mathcal{B}))^m\}$ is a primitive canonical transformation from F to $F \otimes F$ whose graph consists of vertices $x_1^{l_1}, \dots, x_m^{l_m}, y_1^{l_1}, \dots, y_m^{l_m}, y_{m+1}^{l_1}, \dots, y_{m+m}^{l_m}$ and edges $\{x_1, y_1\}, \{x_1, y_{m+1}\}, \dots, \{x_m, y_m\}, \{x_m, y_{2m}\}$.

$\mathbf{k}_F = \{\mathbf{k}_{F(\vec{A})} \mid \vec{A} \in (Ob(\mathcal{B}))^m\}$ is a primitive canonical transformation from F to I with the graph whose vertices are $x_1^{l_1}, \dots, x_m^{l_m}, g_1, \dots, g_m$ and whose edges are $\{x_1, g_1\}, \dots, \{x_m, g_m\}$.

$\epsilon_{F,G} = \{\epsilon_{F(\vec{A}), G(\vec{B})} \mid \vec{A} \in (Ob(\mathcal{B}))^m, \vec{B} \in (Ob(\mathcal{B}))^n\}$ is a primitive canonical transformation from $F \otimes (F \rightarrow G)$ to G with the graph whose vertices are $x_1^{l_1}, \dots, x_m^{l_m}, x_{m+1}^{-l_1}, \dots, x_{m+m}^{-l_m}, x_{2m+1}^{l_{m+1}}, \dots, x_{2m+n}^{l_{m+n}}, y_1^{l_{m+1}}, \dots, y_n^{l_{m+n}}$ and whose edges are $\{x_1, x_{m+1}\}, \dots, \{x_m, x_{2m}\}, \{x_{2m+1}, y_1\}, \dots, \{x_{2m+n}, y_n\}$.

Analogously, we define the primitive canonical transformations δ_F^i from F to $F \otimes I$, $\mathbf{b}_{F,G,H}^{\rightarrow}$ from $F \otimes (G \otimes H)$ to $(F \otimes G) \otimes H$, $\mathbf{b}_{F,G,H}^{\leftarrow}$ from $(F \otimes G) \otimes H$ to $F \otimes (G \otimes H)$, $\mathbf{l}_F$ from O to F , $\mathbf{l}_{F,G}^1$ from F to $F \oplus G$, $\mathbf{l}_{F,G}^2$ from G to $F \oplus G$, $\mathbf{m}_F$ from $F \oplus F$ to F and $\boldsymbol{\eta}_{F,G}$ from G to $F \rightarrow (F \otimes G)$ with corresponding graphs. It is not difficult to show that every primitive canonical transformation is g-dinatural with respect to the associated graph.

Next we define the following operations on canonical transformations.

If $\alpha = \{\alpha(A_1, \dots, A_{k_\Gamma}) \mid A_1, \dots, A_{k_\Gamma} \in Ob(\mathcal{B})\}$ is a canonical transformation from F to G with the graph Γ , then for $l \geq 1$

$$\alpha^{i_1, \dots, i_l} = \{\alpha(A_1, \dots, A_{k_\Gamma}) \mid A_1, \dots, A_{k_\Gamma} \in Ob(\mathcal{B}), A_{i_1} = A_{i_2} = \dots = A_{i_l}\}$$


is a canonical transformation from F to G with the graph obtained from Γ by addition of edges between the vertices from the components $i_1, \dots, i_l$ in order to obtain one component of the new graph. We call $\alpha^{i_1, \dots, i_l}$ a *subtransformation* of α . It is easy to verify that if α and β are canonical transformation from F to G and if $\beta \subset \alpha$, then β is a subtransformation of α . Also, if a canonical transformation is g-dinatural, then each of its subtransformations is g-dinatural, too.

If $\alpha = \{\alpha(A_1, \dots, A_{k_\Phi}) \mid A_1, \dots, A_{k_\Phi} \in Ob(\mathcal{B})\}$ and $\beta = \{\beta(B_1, \dots, B_{k_\Psi}) \mid B_1, \dots, B_{k_\Psi} \in Ob(\mathcal{B})\}$ are two canonical transformations from F to G and from H to J respectively, then


$$\begin{aligned} \alpha \otimes \beta &= \{\alpha(A_1, \dots, A_{k_\Phi}) \times \beta(B_1, \dots, B_{k_\Psi}) \mid A_1, \dots, B_{k_\Psi} \in Ob(\mathcal{B})\}, \\ \alpha \oplus \beta &= \{\alpha(A_1, \dots, A_{k_\Phi}) + \beta(B_1, \dots, B_{k_\Psi}) \mid A_1, \dots, B_{k_\Psi} \in Ob(\mathcal{B})\}, \\ \alpha \rightarrow \beta &= \{\alpha(A_1, \dots, A_{k_\Phi}) \rightarrow \beta(B_1, \dots, B_{k_\Psi}) \mid A_1, \dots, B_{k_\Psi} \in Ob(\mathcal{B})\} \end{aligned}$$

are canonical transformations from $F \otimes H$ to $G \otimes J$, from $F \oplus H$ to $G \oplus J$, and from $G \rightarrow H$ to $F \rightarrow J$ respectively. If Φ is the graph of α and Ψ is the graph of β , then the graphs of $\alpha \otimes \beta$, $\alpha \oplus \beta$ and $\alpha \rightarrow \beta$ are obtained as disjoint unions of Φ and Ψ , where in the last case, Φ occurs inverted. We denote these graphs by $\Phi \otimes \Psi$, $\Phi \oplus \Psi$ and $\Phi \rightarrow \Psi$ respectively.

EXAMPLE 3.8. Let Φ be the graph on left-hand side and Ψ the graph on right-hand side of the picture below.



Then $\Phi \otimes \Psi$ and $\Phi \oplus \Psi$ are identical and given by the diagram on left-hand side and $\Phi \rightarrow \Psi$ is given by the diagram on right-hand side below.



Moreover, canonical transformations are closed under composition defined as in Section 2; i.e. if α is a canonical transformation from F to G with the graph Φ and β is a canonical transformation from G to H with the graph Ψ , then $\beta\alpha$ defined as in Section 2, is a canonical transformation with the graph $\Psi\Phi$.

It is easy to verify that $\mathcal{F}_{\mathcal{B}}$ is a category with the identity morphism for F being $\mathbf{1}_F$ and the composition of α and β being $\beta\alpha$ defined as above. We leave the details about the structure of this category for another occasion.


Our aim is to show that all the morphisms from $\mathcal{F}_{\mathcal{B}}$ are g-dinatural transformations. It is easy to see that the only obstacle for this is the composition of canonical transformations. To show that composition is now harmless too, we use the results from Section 2 and the procedure of cut elimination in an adequate sequent system.

The following example shows that the results from Section 2 are not sufficient for our aims before a further analysis of properties peculiar to bicartesian closed categories.

EXAMPLE 3.9. Let α be the canonical transformation obtained from the following composition of canonical transformations (from now on, we associate compositions to the right)

$$(\epsilon_{1,1} \otimes \mathbf{1}_{1 \rightarrow 1}) \mathbf{b}_{1, (1 \rightarrow 1), (1 \rightarrow 1)}^{\rightarrow} (\epsilon_{1,1} \otimes \mathbf{1}_{(1 \rightarrow 1) \otimes (1 \rightarrow 1)}) \mathbf{b}_{1, 1 \rightarrow 1, (1 \rightarrow 1) \otimes (1 \rightarrow 1)}^{\rightarrow} (\mathbf{1}_1 \otimes (\mathbf{1}_{1 \rightarrow 1} \otimes \mathbf{w}_{1 \rightarrow 1})) (\mathbf{1}_1 \otimes \mathbf{w}_{1 \rightarrow 1})$$

and let β be $\epsilon_{1,1}$. From the facts that the primitive canonical transformations are g-dinatural, that $\otimes$ preserves g-dinaturality, and from Theorem 2.2, it follows that α and β are g-dinatural transformations whose amalgamation of graphs is given by the following diagram.



Since an alternating loop occurs in this amalgamation, by Theorem 2.2. there is a composition of g-dinatural transformations with such graphs, which is not g-dinatural. Of course, it doesn't mean that $\beta\alpha$ is not g-dinatural. However, each element of $\beta\alpha$ is in the composition of canonical transformations

$$\alpha_1 = \mathbf{b}_{1 \otimes (1 \rightarrow 1), (1 \rightarrow 1), (1 \rightarrow 1)}^{\rightarrow} \mathbf{b}_{1, 1 \rightarrow 1, (1 \rightarrow 1) \otimes (1 \rightarrow 1)}^{\rightarrow} (\mathbf{1}_1 \otimes (\mathbf{1}_{1 \rightarrow 1} \otimes \mathbf{w}_{1 \rightarrow 1})) (\mathbf{1}_1 \otimes \mathbf{w}_{1 \rightarrow 1})$$

and

$$\beta_1 = \epsilon_{1,1} (\epsilon_{1,1} \otimes \mathbf{1}_{1 \rightarrow 1}) (\epsilon_{1,1} \otimes \mathbf{1}_{(1 \rightarrow 1) \otimes (1 \rightarrow 1)})$$

which in turn gives that $\beta\alpha$ is a subtransformation of $\beta_1\alpha_1$. The g-dinaturality of β_1 , and of $\beta_1\alpha_1$ too, follows from Theorem 2.2. Hence, from these two facts it follows that $\beta\alpha$ is g-dinatural. In the sequel, we generalize the idea from the example above to the case of an arbitrary composition of canonical transformations. For this purpose we need the following definitions.

Let α be a canonical transformation. Denote by $C(\alpha)$ the set of canonical transformations defined inductively by

- $\alpha \in C(\alpha)$,
- if $\beta \in C(\alpha)$ and $F \in Ob(\mathcal{F}_{\mathcal{B}})$, then $(\beta \otimes \mathbf{1}_F)$ and $(\mathbf{1}_F \otimes \beta)$ are in $C(\alpha)$.

Let $\xi_{F,G,H}$ from $F \otimes (G \oplus H)$ to $(F \otimes G) \oplus (F \otimes H)$ be the following canonical transformation.

$$\epsilon_{F, (F \otimes G) \oplus (F \otimes H)} (\mathbf{1}_F \otimes \mathbf{m}_{F \rightarrow ((F \otimes G) \oplus (F \otimes H))}) ((\mathbf{1}_F \rightarrow \mathbf{l}_{F \otimes G, F \otimes H}^1) \eta_{F,G} + ((\mathbf{1}_F \rightarrow \mathbf{l}_{F \otimes G, F \otimes H}^2) \eta_{F,H}))$$

Next we define the set $Constr(\mathcal{B})$ of *constructible* canonical transformations. This name comes from the analogous notion from [8].

1. Primitive canonical transformations are in $\text{Constr}(\mathcal{B})$.
2. If α from F to G is in $C(\beta)$ for β be among $\mathbf{b}_{T,S,R}^{\leftarrow}$, $\mathbf{b}_{T,S,R}^{\rightarrow}$, $\mathbf{c}_{T,S}$, $\mathbf{w}_T$, $\mathbf{k}_T$, $\boldsymbol{\delta}_T$, $\boldsymbol{\delta}_T^i$ for some $T, S, R \in \text{Ob}(\mathcal{F}_{\mathcal{B}})$, and γ from G to H is in $\text{Constr}(\mathcal{B})$, then $\gamma\alpha$ is in $\text{Constr}(\mathcal{B})$.
3. If α and β are in $\text{Constr}(\mathcal{B})$ then $\alpha \otimes \beta$ is in $\text{Constr}(\mathcal{B})$.
4. If α from $F \otimes G$ to H and β from $J \otimes G$ to H are in $\text{Constr}(\mathcal{B})$, then $\mathbf{m}_H(\alpha \oplus \beta)(\mathbf{c}_{G,F} \oplus \mathbf{c}_{G,J})\xi_{F,J,G}\mathbf{c}_{F \oplus J,G}$ is in $\text{Constr}(\mathcal{B})$.
5. If α from F to G is in $\text{Constr}(\mathcal{B})$ then $\mathbf{l}_{G,H}^1\alpha$ and $\mathbf{l}_{G,H}^2\alpha$ are in $\text{Constr}(\mathcal{B})$.
6. If α from $F \otimes G$ to H is in $\text{Constr}(\mathcal{B})$ then $(\mathbf{1}_F \rightarrow \alpha)\boldsymbol{\eta}_{F,G}$ is in $\text{Constr}(\mathcal{B})$.
7. If α from F to G and β from $H \otimes J$ to T are in $\text{Constr}(\mathcal{B})$ then $\beta((\boldsymbol{\varepsilon}_{G,H}(\alpha \otimes \mathbf{1}_{G \rightarrow H})) \otimes \mathbf{1}_J)$ is in $\text{Constr}(\mathcal{B})$.

LEMMA 3.13. *Each constructible canonical transformation is g-dinatural with respect to its own graph.*

PROOF. It is easy to verify that the primitive canonical transformations are g-dinatural. (This follows from the equations $(\boldsymbol{\delta})$, $(\mathbf{b})$, $(\mathbf{c})$, $(\mathbf{w})$, $(\mathbf{k})$, $(\mathbf{l}^1)$, $(\mathbf{l}^2)$, $(\mathbf{l})$, $(\mathbf{m})$, $(\boldsymbol{\varepsilon}1)$, $(\boldsymbol{\eta}1)$, $(\boldsymbol{\varepsilon}2)$ and $(\boldsymbol{\eta}2)$.) Also it is easy to see that if α and β are g-dinatural, then such are $\alpha \otimes \beta$, $\alpha \oplus \beta$ and $\alpha \rightarrow \beta$, too. For the rest, we rely on Theorem 2.2.

4 A Category-like Sequent System for Intuitionistic Propositional Logic

In this section we carry out a cut elimination procedure in an auxiliary sequent system for intuitionistic propositional logic, which will help us in dealing with the dinaturality of bicartesian closed canonical transformations.

This sequent system, which we call, $\mathcal{J}$ is introduced as follows. Let $\mathcal{F}$ be generated from a countable set $\mathcal{L}$, whose members we call *propositional letters*, with the constants $\top$ and $\perp$ and the binary connectives $\wedge$, $\vee$ and $\rightarrow$. We call the members of $\mathcal{F}$ *formulae*, and use the schematic letters $A, B, C, \dots, A_1, \dots$ for them. *Sequents* of $\mathcal{J}$ are of the form $A \vdash B$ for A and B in $\mathcal{F}$. We call A in $A \vdash B$ the *antecedent*, and B the *consequent* of the sequent. In order to introduce the rules of inference of $\mathcal{J}$ we need the following auxiliary notion of $\wedge$ -context, which corresponds to the notion of (poly)functor in categories. A $\wedge$ -context is defined inductively as follows:

- 1° the symbol $\square$ is a $\wedge$ -context,
- 2° if G is a $\wedge$ -context and $A \in \mathcal{F}$, then $(G \wedge A)$ and $(A \wedge G)$ are $\wedge$ -contexts,
- 3° if G and H are $\wedge$ -contexts, then $(G \wedge H)$ is a $\wedge$ -context.

For a $\wedge$ -context F we say that it is a $\wedge_1$ -context if the symbol $\square$ occurs in F exactly once. For G a $\wedge$ -context and $A \in \mathcal{F}$, we obtain $G(A)$ by substituting A for $\square$ in G , e.g., if $F \equiv (B \wedge \square) \wedge C$, then $F(A) = (B \wedge A) \wedge C$.

The *axioms* of $\mathcal{J}$ are

$$a_A : A \vdash A, \quad \Pi_A : \perp \vdash A, \quad \text{for every } A \in \mathcal{F},$$

The *structural rules* of $\mathcal{J}$ are

$$\begin{array}{ll}
(\beta_F^-) \frac{F(A \wedge (B \wedge C)) \vdash D}{F((A \wedge B) \wedge C) \vdash D} & (\beta_F^+) \frac{F((A \wedge B) \wedge C) \vdash D}{F(A \wedge (B \wedge C)) \vdash D} \\
(\gamma_F) \frac{F(A \wedge B) \vdash C}{F(B \wedge A) \vdash C} & \\
(\omega_F) \frac{F(A \wedge A) \vdash B}{F(A) \vdash B} & (\theta_F^A) \frac{F(\top) \vdash B}{F(A) \vdash B} \\
(\tau_F) \frac{F(A) \vdash B}{F(A \wedge \top) \vdash B} & (\tau_F^i) \frac{F(A \wedge \top) \vdash B}{F(A) \vdash B} \\
(\odot) \frac{A \vdash B \quad G(B) \vdash C}{G(A) \vdash C}, &
\end{array}$$

where F is a $\wedge_1$ context and G is a $\wedge$ context. The last rule is called *mix* and we refer to it by $(\odot)$ when the context G is irrelevant.

The *rules for connectives* are

$$\begin{array}{ll}
(\wedge) \frac{A \vdash C \quad B \vdash D}{A \wedge B \vdash C \wedge D} & \\
(\diamond) \frac{A \wedge C \vdash D \quad B \wedge C \vdash D}{(A \vee B) \wedge C \vdash D} & (+c) \frac{A \vdash B}{A \vdash B \vee C} \quad (b+) \frac{A \vdash C}{A \vdash B \vee C} \\
(*) \frac{A \wedge B \vdash C}{B \vdash A \rightarrow C} & (\triangleright) \frac{A \vdash B \quad C \wedge D \vdash E}{(A \wedge (B \rightarrow C)) \wedge D \vdash E}
\end{array}$$

A *proof* of a sequent $A \vdash B$ in $\mathcal{J}$ is a binary tree with sequents in its nodes, such that $A \vdash B$ is in the root, axioms are in the leaves and consecutive nodes are connected by some of the inference rules above.

It is not difficult to see that the underlying logic of $\mathcal{J}$ is intuitionistic propositional logic. The differences between $\mathcal{J}$ and Gentzen's system LJ introduced in [6] are that in $\mathcal{J}$ we have just one meta-logical symbol ($\vdash$) in the sequents: we omit Gentzen's commas in the antecedents, whose role is now covered by the logical connective $\wedge$. We can't have empty either the antecedent or the consequent of a sequent in $\mathcal{L}$. The logical constant $\top$ serves to fill gaps in antecedents. These discrepancies between $\mathcal{J}$ and LJ arise because in $\mathcal{J}$ we want antecedents and consequents of sequents to be of the same sort (namely members of $\mathcal{F}$) and this enables us to look at an $\mathcal{J}$ sequent as an arrow with the source being the antecedent and the target the consequent of the sequent.

Our $(\wedge)$ is a rule of simultaneous introduction of the connective $\wedge$ on the both sides of a sequent: there is no a counterpart for this rule in LJ . This difference is not categorially motivated though it emphasizes functoriality of the connective $\wedge$. We also believe that $\mathcal{J}$ completely separates structural rules from the rules for connectives. On the other hand, the LJ rules $\&$ -IS and $\&$ -IA (see 1.22. of [6]) have hidden interchanges, contractions and thinnings.

Since we prove the cut-elimination theorem through elimination of *mix*, as Gentzen did too, we have postulated the mix rule $(\odot)$ as primitive. However, this mix is something different from Gentzen's mix. It is liberal in the sense that the $\wedge$ -context G in $(\odot)$ need not to capture all factors B (see the definition below) as arguments in $G(B)$. This means that the formula B may be a factor of A in Step 2° of the construction of the $\wedge$ -context G ; i.e. mix need not to “swallow” all the occurrences of B in $G(B)$. There are no categorial reasons to prefer cut to such a mix. In both cases, we don't have categorial composition of arrows corresponding to both premises of the rule, but a more involved

composition of the right premise with an image of the left premise under the functor corresponding to a $\wedge$ -context. The only difference is that in the case of cut this is always a $\wedge_1$ -context.

An advantage of $\mathcal{J}$ is that its proofs can be easily coded. For example the proof

$$\frac{\frac{p \vdash p \quad \frac{q \vdash q}{q \wedge \top \vdash q}}{(p \wedge (p \rightarrow q)) \wedge \top \vdash q}}{p \wedge (p \rightarrow q) \vdash q}$$

is coded by

$$\tau_{\square}^i(a_p \triangleright \tau_{\square} a_q)$$

This fact helps when we want to postulate equalities that should hold between the proofs of $\mathcal{J}$.

For G a $\wedge$ -context and π a proof, we denote by $G(\pi)$ the proof coded by the term obtained from G after the substitution a_A for every A and the code of π for every $\square$ in G .

For the proof of the main result of this section we need the following notions of *degree* and *rank*. The degree of a formula is the number of logical connectives in it. However, because of the categorially motivated elimination of the comma, the symbol $\wedge$ plays a double role and in order to define rank, we define as follows a set of *factors* of A , for every $A \in \mathcal{F}$:

1° A is a factor of A ,

2° if A is of the form $A_1 \wedge A_2$ then every factor of A_1 or A_2 is a factor of A .

Now, we introduce (in the style of Došen) an auxiliary indexing of consequents and factors of antecedents in a mixless proof of $\mathcal{J}$ which will help us in defining the rank of an occurrence of a formula in such a proof. First we index all the consequents and all the factors of antecedents of axioms by 1 and inductively proceed as follows. In all the structural rules and the rule $(\triangleright)$ the index of the consequent in the conclusion is increased by 1. In $(\diamond)$ the index of the consequent in the conclusion is the maximum of the two indices of consequents of both premises increased by 1. In $(\wedge)$, $(+c)$, $(c+)$ and $(*)$ the index of the consequent in the conclusion is 1. Every factor of the antecedent preserved by a rule has the index increased by 1, and all the factors introduced by the rule have index 1 in the conclusion. In (ω_F) the occurrence of A in the conclusion is indexed by the maximum of the indices of distinguished A 's in the premise, increased by 1. In the example of the proof given above this indexing looks like

$$\frac{\frac{p^1 \vdash p^1 \quad \frac{q^1 \vdash q^1}{(q^2 \wedge \top^1)^1 \vdash q^2}}{((p^2 \wedge (p \rightarrow q)^1)^1 \wedge \top^2)^1 \vdash q^3}}{(p^3 \wedge (p \rightarrow q)^2)^2 \vdash q^4}$$

Then the rank of an occurrence of a formula in a mixless proof is given by its index.

The following theorem corresponds to Gentzen's *Hauptsatz* of [6].

THEOREM 4.3. *Every proof in $\mathcal{J}$ can be transformed into a proof of the same root-sequent with no applications of the rule $(\bigcirc)$.*

PROOF. As in the standard cut-elimination procedure it is enough to consider a proof π whose last rule is $(\bigodot)$ for a $\wedge$ -context G , and there is no more application of $(\bigcirc)$ in π . So let our proof be of the form

$$\frac{\frac{\pi_1}{A \vdash B} \quad \frac{\pi_2}{G(B) \vdash C}}{G(A) \vdash C} \bigodot$$

with π_1 and π_2 mixless. Then we define the *degree of this proof* as the degree of B and the *rank of this proof* as the sum of the *left rank*, i.e. the rank of the occurrence of B in the left premise of $\odot$, in the subproof π_1 , and the *right rank*, i.e. the maximum of all ranks of distinguished factors B in the right premise of $\odot$ in the subproof π_2 . Then we prove our theorem by induction on the lexicographically ordered pairs $\langle d, r \rangle$ for the degree d and the rank r of the proof.

1. $r = 2$

1.1. π_1 or π_2 are axioms

1.1.1. Suppose π is of the form

$$\frac{a_B : B \vdash B \quad \begin{array}{c} \pi_2 \\ G(B) \vdash C \end{array}}{G(B) \vdash C}$$

Then this proof is transformed into the proof

$$\begin{array}{c} \pi_2 \\ G(B) \vdash C \end{array}$$

which is mixless.

1.1.2. If π is of the form

$$\frac{\begin{array}{c} \pi_1 \\ A \vdash B \end{array} \quad a_{G(B)} : G(B) \vdash G(B)}{G(A) \vdash G(B)}$$

Then this proof is transformed into the proof

$$\begin{array}{c} G(\pi_1) \\ G(A) \vdash G(B) \end{array}$$

which is of course mixless.

1.1.3. If π is of the form

$$\frac{\Pi_B : \perp \vdash B \quad \begin{array}{c} \pi_2 \\ G(B) \vdash C \end{array}}{G(\perp) \vdash C}$$

Then this proof is transformed into the proof of the form

$$\frac{\frac{\Pi_C : \perp \vdash C}{\dots} \quad (\tau), (\gamma), (\theta)}{G(\perp) \vdash C}$$

1.1.4. Finally, if π is of the form

$$\frac{\begin{array}{c} \pi_1 \\ A \vdash \perp \end{array} \quad \Pi_C : \perp \vdash C}{A \vdash C}$$

Then, since the left rank of this proof is 1, A must be $\perp$ and π is transformed into $\Pi_C : \perp \vdash C$.

1.2. π_1 ends with $(\wedge)$

Suppose π is of the form

$$\frac{\frac{\pi'_1}{A_1 \vdash B_1} \quad \frac{\pi''_1}{A_2 \vdash B_2}}{A_1 \wedge A_2 \vdash B_1 \wedge B_2} \wedge \quad \frac{\pi_2}{G(B_1 \wedge B_2) \vdash C} \quad \textcircled{G}$$

$$\frac{}{G(A_1 \wedge A_2) \vdash C}$$

Then this proof is transformed into the proof

$$\frac{\pi'_1}{A_1 \vdash B_1} \quad \frac{\frac{\pi''_1}{A_2 \vdash B_2} \quad \frac{\pi_2}{G(B_1 \wedge B_2) \vdash C}}{G(B_1 \wedge A_2) \vdash C} \quad \textcircled{}$$

$$\frac{}{G(A_1 \wedge A_2) \vdash C} \quad \textcircled{}$$

where both applications of $(\textcircled{})$ have lower degree.

1.3. π_1 ends with $(*)$

1.3.1. π_2 ends with (θ)

Suppose π is of the form

$$\frac{\frac{\pi'_1}{B_1 \wedge A \vdash B_2}}{A \vdash B_1 \rightarrow B_2} * \quad \frac{\frac{\pi'_2}{F(\top) \vdash C}}{G(B_1 \rightarrow B_2) \vdash C} \theta_F^{G_1(B_1 \rightarrow B_2)}$$

$$\frac{}{G(A) \vdash C}$$

Then this proof is transformed into the proof

$$\frac{\pi'_2}{F(\top) \vdash C} \theta_F^{G_1(A)}$$

$$\frac{}{G(A) \vdash C}$$

1.3.2. π_2 ends with $(\triangleright)$

Suppose π is of the form

$$\frac{\frac{\pi'_1}{B_1 \wedge A \vdash B_2}}{A \vdash B_1 \rightarrow B_2} * \quad \frac{\frac{\pi'_2}{D \vdash B_1} \quad \frac{\pi''_2}{B_2 \wedge E \vdash C}}{(D \wedge (B_1 \rightarrow B_2)) \wedge E \vdash C} \triangleright$$

$$\frac{}{(D \wedge A) \wedge E \vdash C}$$

Then this proof is transformed into the proof

$$\frac{\pi'_2}{D \vdash B_1} \quad \frac{\frac{\pi'_1}{B_1 \wedge A \vdash B_2} \quad \frac{\pi''_2}{B_2 \wedge E \vdash C}}{(B_1 \wedge A) \wedge E \vdash C} \quad \textcircled{}$$

$$\frac{}{(D \wedge A) \wedge E \vdash C} \quad \textcircled{}$$

with both applications of $(\bigcirc)$ of the lower degree.

1.4. π_1 ends with $(+_{B_2})$, or analogously with $(_{B_1}+)$

1.4.1. π_2 ends with (θ) is analogous to 1.3.1.

1.4.2. π_2 ends with $(\diamond)$

Suppose π is of the form

$$\frac{\frac{\pi'_1}{A \vdash B_1} \quad \frac{\pi'_2 \quad \pi''_2}{\frac{B_1 \wedge D \vdash C \quad B_2 \wedge D \vdash C}{(B_1 \vee B_2) \wedge D \vdash C} \diamond} \quad +_{B_2} \quad \frac{A \vdash B_1 \vee B_2}{A \wedge D \vdash C}$$

Then this proof is transformed into the proof

$$\frac{\pi'_1 \quad \pi'_2}{\frac{A \vdash B_1 \quad B_1 \wedge D \vdash C}{A \wedge D \vdash C} \bigcirc} \bigcirc$$

with the smaller degree.

2. $r > 2$

2.1. the right rank is > 1

2.1.1. π_2 ends with a structural rule (σ) , i.e., π is of the form

$$\frac{\pi_1 \quad \frac{\pi_2}{\frac{G_1(B) \vdash C}{G(B) \vdash C} \sigma}}{A \vdash B \quad \frac{G_1(B) \vdash C}{G(B) \vdash C} \sigma} \sigma \quad \frac{A \vdash B \quad G(B) \vdash C}{G(A) \vdash C}$$

2.1.1.1. If all the distinguished B 's in the right premise of $(\mathcal{G})$ in π have indices greater than 1 (by 2.1, at least one such B must occur) then this proof is transformed into the proof

$$\frac{\pi_1 \quad \pi_2}{\frac{A \vdash B \quad G_1(B) \vdash C}{G_1(A) \vdash C} \mathcal{G}_1} \sigma \quad \frac{G_1(A) \vdash C}{G(A) \vdash C}$$

whose subproof ending with $(\mathcal{G}_1)$ has the rank lower by 1.

2.1.1.2. If one of the distinguished B 's in the right premise of $(\mathcal{G})$ in π is indexed by 1 (note that except for (θ) , in the conclusion of a structural rule, every formula has at most one occurrence indexed by 1), then π is transformed into the proof

$$\frac{\pi_1 \quad \pi_2}{\frac{A \vdash B \quad G_1(B) \vdash C}{G_1(A) \vdash C} \mathcal{G}_1} \sigma \quad \frac{A \vdash B \quad \frac{G_1(A) \vdash C}{F(B) \vdash C} \sigma}{G(A) \vdash C} \mathcal{F}$$

for a $\wedge_1$ -context F (except when σ is an application of (θ) in which case F is a $\wedge$ -context) such that $F(A) \equiv G(A)$. In this proof, the subproof ending with the upper mix has the rank decreased by 1, and the right rank of the lower mix remains equal to 1 after the elimination of the upper mix.

2.1.2. π_2 ends with $(\wedge)$

Suppose π is of the form

$$\frac{\pi_1 \quad \frac{\pi'_2 \quad G_1(B) \vdash C_1 \quad \pi''_2 \quad G_2(B) \vdash C_2}{G(B) \vdash C_1 \wedge C_2} \wedge}{A \vdash B \quad G(A) \vdash C_1 \wedge C_2} \wedge$$

Then this proof is transformed into the proof

$$\frac{\frac{\pi_1 \quad A \vdash B}{G_1(A) \vdash C_1} \textcircled{G_1} \quad \frac{\pi_1 \quad A \vdash B \quad \pi''_2 \quad G_2(B) \vdash C_2}{G_2(A) \vdash C_2} \textcircled{G_2}}{G(A) \vdash C_1 \wedge C_2} \wedge$$

in which both subproofs ending with $\textcircled{G_1}$ and $\textcircled{G_2}$ are of the lower ranks. There is also a simplified variant of 2.1.2 with no distinguished B 's in the antecedent of a premise of the rule $(\wedge)$.

In all the cases below, the subproofs of the reduced proofs ending with the applications of $(\bigcirc)$, have a smaller rank than π .

2.1.3. π_2 ends with $(*)$

Suppose π is of the form

$$\frac{\pi_1 \quad \frac{\pi'_2 \quad C_1 \wedge G(B) \vdash C_2}{G(B) \vdash C_1 \rightarrow C_2} *}{A \vdash B \quad G(A) \vdash C_1 \rightarrow C_2} *$$

Then this proof is transformed into the proof

$$\frac{\pi_1 \quad A \vdash B \quad \pi_2 \quad C_1 \wedge G(B) \vdash C_2}{C_1 \wedge G(A) \vdash C_2} \bigcirc \quad \frac{C_1 \wedge G(A) \vdash C_2}{G(A) \vdash C_1 \rightarrow C_2} *$$

2.1.4. π_2 ends with $(\triangleright)$

2.1.4.1. Suppose that π is of the form

$$\frac{\pi_1 \quad A \vdash B \quad \frac{\pi'_2 \quad G_1(B) \vdash B_1 \quad \pi''_2 \quad B_2 \wedge G_2(B) \vdash C}{(G_1(B) \wedge B) \wedge G_2(B) \vdash C}}{(G_1(A) \wedge A) \wedge G_2(A) \vdash C} \triangleright$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\begin{array}{c}
\pi_1 \quad \pi'_2 \\
\frac{A \vdash B \quad G_1(B) \vdash B_1}{G_1(A) \vdash B_1} \circ
\end{array}
\quad
\begin{array}{c}
\pi_1 \quad \pi''_2 \\
\frac{A \vdash B \quad B_2 \wedge G_2(B) \vdash C}{B_2 \wedge G_2(A) \vdash C} \circ
\end{array} \\
\frac{\pi_1 \quad \frac{A \vdash B \quad G_1(A) \vdash B_1}{G_1(A) \vdash B_1} \quad \frac{B_2 \wedge G_2(A) \vdash C}{B_2 \wedge G_2(A) \vdash C}}{(G_1(A) \wedge B) \wedge G_2(A) \vdash C} \triangleright \\
\frac{A \vdash B \quad (G_1(A) \wedge B) \wedge G_2(A) \vdash C}{(G_1(A) \wedge A) \wedge G_2(A) \vdash C} \circ
\end{array}$$

2.1.4.2. Suppose that π is of the form

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{B_1 \vdash B_2 \quad B_3 \wedge G_1(B) \vdash C}{B \wedge G_1(B) \vdash C} \triangleright}{A \wedge G_1(A) \vdash C}
\end{array}$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{B_1 \vdash B_2 \quad \frac{A \vdash B \quad B_3 \wedge G_1(B) \vdash C}{B_3 \wedge G_1(A) \vdash C} \circ}{B \wedge G_1(A) \vdash C} \triangleright}{A \wedge G_1(A) \vdash C} \circ
\end{array}$$

2.1.4.3. Suppose that π is of the form

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{G_1(B) \vdash D \quad E \wedge G_2(B) \vdash C}{(G_1(B) \wedge (D \rightarrow E)) \wedge G_2(B) \vdash C} \triangleright}{(G_1(A) \wedge (D \rightarrow E)) \wedge G_2(A) \vdash C}
\end{array}$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{G_1(B) \vdash D}{G_1(A) \vdash D} \circ \quad \frac{A \vdash B \quad E \wedge G_2(B)}{E \wedge G_2(A) \vdash C} \circ}{(G_1(A) \wedge (D \rightarrow E)) \wedge G_2(A) \vdash C} \triangleright
\end{array}$$

There are also simplified variants of 2.1.4.1. and 2.1.4.3. with no distinguished B 's in G_1 or G_2 which we won't discuss here separately.

2.1.5. π_2 ends with ($\diamond$)

2.1.5.1. Suppose that π is of the form

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{B_1 \wedge G_1(B) \vdash C \quad B_2 \wedge G_1(B) \vdash C}{B \wedge G_1(B) \vdash C} \diamond}{A \wedge G_1(A) \vdash C}
\end{array}$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\begin{array}{c}
\pi_1 \quad \pi'_2 \\
\frac{A \vdash B \quad B_1 \wedge G_1(B) \vdash C}{B_1 \wedge G_1(A) \vdash C} \bigcirc
\end{array}
\quad
\begin{array}{c}
\pi_1 \quad \pi''_2 \\
\frac{A \vdash B \quad B_2 \wedge G_1(B) \vdash C}{B_2 \wedge G_1(A) \vdash C} \bigcirc
\end{array} \\
\frac{\pi_1 \quad \frac{B_1 \wedge G_1(A) \vdash C \quad B_2 \wedge G_1(A) \vdash C}{B \wedge G_1(A) \vdash C} \diamond}{A \vdash B \quad B \wedge G_1(A) \vdash C} \bigcirc \\
\hline
A \wedge G_1(A) \vdash C
\end{array}$$

2.1.5.2. Suppose that π is of the form

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{D_1 \wedge G_1(B) \vdash C \quad D_2 \wedge G_1(B) \vdash C}{(D_1 \vee D_2) \wedge G_1(B) \vdash C} \diamond}{(D_1 \vee D_2) \wedge G_1(A) \vdash C}
\end{array}$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\pi_1 \quad \pi'_2 \quad \pi''_2 \\
\frac{A \vdash B \quad \frac{D_1 \wedge G_1(B) \vdash C}{D_1 \wedge G_1(A) \vdash C} \bigcirc \quad \frac{A \vdash B \quad \frac{D_2 \wedge G_1(B) \vdash C}{B_2 \wedge G_1(A) \vdash C} \bigcirc}{(D_1 \vee D_2) \wedge G_1(A) \vdash C} \diamond
\end{array}$$

2.1.6. π_2 ends with $(+_{C_2})$

Suppose that π is of the form

$$\begin{array}{c}
\pi_2 \\
\frac{\pi_1 \quad \frac{G(B) \vdash C_1}{G(B) \vdash C_1 \vee C_2} +_{C_2}}{G(A) \vdash C_1 \vee C_2}
\end{array}$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\pi_1 \quad \pi_2 \\
\frac{A \vdash B \quad G(B) \vdash C_1}{G(A) \vdash C_1} \bigcirc \\
\frac{G(A) \vdash C_1}{G(A) \vdash C_1 \vee C_2} +_{C_2}
\end{array}$$

The case of (c_{1+}) instead of $(+_{C_2})$ is dealt with analogously.

2.2. The right rank is 1 and the left rank is greater than 1.

If π_2 is the axiom $a_{G(B)}$, then we proceed as in 1.1.2. If π_2 ends with an application of (θ) , then we proceed as in 1.3.1. In all the remaining cases G must be a $\wedge_1$ -context

2.2.1. π_1 ends with a structural rule

Suppose that π is of the form

$$\frac{\frac{\pi'_1}{A_1 \vdash B} \quad \sigma \quad \frac{\pi_2}{G(B) \vdash C}}{G(A) \vdash C}$$

Then this proof is transformed into the proof

$$\frac{\frac{\pi'_1}{A_1 \vdash B} \quad \frac{\pi_2}{G(B) \vdash C}}{G(A_1) \vdash C} \circ \quad \frac{G(A_1) \vdash C}{G(A) \vdash C} \sigma$$

2.2.2. π_1 ends with $(\triangleright)$

Suppose that π is of the form

$$\frac{\frac{\pi'_1}{A_1 \vdash A_2} \quad \frac{\pi''_1}{A_3 \wedge A_4 \vdash B}}{(A_1 \wedge (A_2 \rightarrow A_3)) \wedge A_4 \vdash B} \triangleright \quad \frac{\pi_2}{G(B) \vdash C}}{G((A_1 \wedge (A_2 \rightarrow A_3)) \wedge A_4) \vdash C}$$

Then this proof is transformed into the proof

$$\frac{\frac{\pi'_1}{A_1 \vdash A_2} \quad \frac{\frac{\frac{\pi''_1}{A_3 \wedge A_4 \vdash B} \quad \frac{\pi_2}{G(B) \vdash C}}{G(A_3 \wedge A_4) \vdash C} \circ \quad \frac{\dots}{(\beta), (\gamma)}}{A_3 \wedge D \vdash C} \quad \frac{A_1 \vdash A_2 \quad A_3 \wedge D \vdash C}{(A_1 \wedge (A_2 \rightarrow A_3)) \wedge D \rightarrow C} \triangleright}{\frac{\dots}{(\beta), (\gamma)}}{G((A_1 \wedge (A_2 \rightarrow A_3)) \wedge A_4) \vdash C}$$

2.2.3. Eventually, if π_1 ends with $(\diamond)$ and π is of the form

$$\frac{\frac{\pi'_1}{A_1 \wedge A_3 \vdash B} \quad \frac{\pi''_1}{A_2 \wedge A_3 \vdash B}}{(A_1 \vee A_2) \wedge A_3 \vdash B} \diamond \quad \frac{\pi_2}{G(B) \vdash C}}{G((A_1 \vee A_2) \wedge A_3) \vdash C}$$

Then this proof is transformed into the proof

$$\begin{array}{c}
\frac{\pi'_1 \quad \pi_2}{\frac{A_1 \wedge A_3 \vdash B \quad G(B) \vdash C}{G(A_1 \wedge A_3) \vdash C} \circ} \quad \frac{\pi''_1 \quad \pi_2}{\frac{A_2 \wedge A_3 \vdash B \quad G(B) \vdash C}{G(A_2 \wedge A_3) \vdash C} \circ} \\
\frac{\dots}{A_1 \wedge D \vdash C} (\beta), (\gamma) \quad \frac{\dots}{A_2 \wedge D \vdash C} (\beta), (\gamma) \\
\hline
\frac{(A_1 \vee A_2) \wedge D \vdash C}{\dots} (\beta), (\gamma) \\
\hline
G((A_1 \vee A_2) \wedge D) \vdash C \quad \diamond
\end{array}$$

□

5 The embedding of $\mathcal{J}$ into a free bicartesian closed category

Let **BiCartCl** be the bicartesian closed category freely generated by the set of objects $\mathcal{L}$ used in Section 4. The morphisms of this category can be viewed as equivalence classes of *morphism terms* generated from $\mathbf{1}_A$, δ_A , δ_A^i , $\mathbf{b}_{A,B,C}^{\rightarrow}$, $\mathbf{b}_{A,B,C}^{\leftarrow}$, $\mathbf{c}_{A,B}$, $\mathbf{w}_A$, $\mathbf{k}_A$, $\varepsilon_{A,B}$, $\eta_{A,B}$, $\mathbf{l}_A$, $\mathbf{l}_{A,B}^1$, $\mathbf{l}_{A,B}^2$ and $\mathbf{m}_A$ for some objects A, B, C of **BiCartCl** with the operations $\times$, $+$, $\rightarrow$ and $\circ$, modulo bicartesian closed equations given in Section 1.

Now we define translations from the set of $\mathcal{J}$ -formulae and the set of $\mathcal{J}$ -proofs to $Ob(\mathbf{BiCartCl})$ and the set of morphism terms, respectively. Denote both these translations by t .

Let t be the identity on $\mathcal{L}$ and inductively defined as follows. (In the following definition, $\mathbf{F}$ is a naturally extracted functor from the $\wedge$ -context F , and the indices of special morphisms can be easily reconstructed.)

$$\begin{aligned}
t(\top) &= \mathbf{I}, & t(\perp) &= \mathbf{O} \\
t(A \wedge B) &= t(A) \times t(B), & t(A \vee B) &= t(A) + t(B), & t(A \rightarrow B) &= t(A) \rightarrow t(B), \\
t(a_A) &= \mathbf{1}_{t(A)}, & t(\Pi_A) &= \mathbf{l}_{t(A)}, \\
t(\beta_F^{\rightarrow}(\pi)) &= t(\pi) \circ \mathbf{F}(\mathbf{b}^{\rightarrow}), & t(\beta_F^{\leftarrow}(\pi)) &= t(\pi) \circ \mathbf{F}(\mathbf{b}^{\leftarrow}), \\
t(\gamma_F(\pi)) &= t(\pi) \circ \mathbf{F}(\mathbf{c}), & t(\theta_F^A(\pi)) &= t(\pi) \circ \mathbf{F}(\mathbf{k}_A), \\
t(\omega_F(\pi)) &= t(\pi) \circ \mathbf{F}(\mathbf{w}), & t(\tau_F^i(\pi)) &= t(\pi) \circ \mathbf{F}(\delta^i), \\
t(\tau_F(\pi)) &= t(\pi) \circ \mathbf{F}(\delta), & & \\
t(\pi_2 \odot \pi_1) &= t(\pi_2) \circ \mathbf{G}(t(\pi_1)), & & \\
t(\pi_1 \wedge \pi_2) &= t(\pi_1) \times t(\pi_2), & & \\
t(\pi_1 \diamond \pi_2) &= \mathbf{m} \circ (t(\pi_1) + t(\pi_2)) \circ (\mathbf{c} + \mathbf{c}) \circ \xi \circ \mathbf{c}, & & \\
t(\pi^{+C}) &= \mathbf{l}_{-,C}^1 \circ t(\pi), & t(\pi^{C+}) &= \mathbf{l}_{C,-}^2 \circ t(\pi), \\
t(\pi^*) &= (\mathbf{1} \rightarrow t(\pi)) \circ \eta, & & \\
t(\pi_1 \triangleright \pi_2) &= t(\pi_2) \circ ((\varepsilon \circ (t(\pi_1) \times \mathbf{1})) \times \mathbf{1}).
\end{aligned}$$

The translation t' that is inverse to t on the set $Ob(\mathbf{BiCartCl})$ is defined on the set of morphism

terms as follows. (Here we write A' instead of $t'(A)$ and f' instead of $t'(f)$.)

$$\begin{aligned}
t'(\mathbf{1}_A) &= a_{A'}, & t'(\mathbf{l}_A) &= \Pi_{A'}, \\
t'(\mathbf{b}_{A,B,C}^\rightarrow) &= \beta_{\Box}^\rightarrow a_{A' \wedge (B' \wedge C')}, & t'(\mathbf{b}_{A,B,C}^\leftarrow) &= \beta_{\Box}^\leftarrow a_{(A' \wedge B') \wedge C'}, \\
t'(\mathbf{c}_{A,B}) &= \gamma_{\Box} a_{B' \wedge A'}, & & \\
t'(\mathbf{w}_A) &= \omega_{\Box} a_{A' \wedge A'}, & t'(\mathbf{k}_A) &= \theta_{\Box}^{A'} a_{\top}, \\
t'(\delta_A) &= \tau_{\Box} a_{A'}, & t'(\delta_A^i) &= \tau_{\Box}^i a_{A' \wedge \top}, \\
t'(\mathbf{l}_{A,B}^1) &= (a_{A'})^{+B'}, & t'(\mathbf{l}_{A,B}^2) &= (a_{B'})^{A'+}, \\
t'(\mathbf{m}_A) &= \tau_{\Box}^i(\tau_{\Box} a_{A'} \diamond \tau_{\Box} a_{a'}), & & \\
t'(\epsilon_{A,B}) &= \tau_{\Box}^i(a_{A'} \triangleright \tau_{\Box} a_{B'}), & t'(\eta_{A,B}) &= (a_{A' \wedge B'})^*, \\
t'(f \times g) &= f' \wedge g', & t'(f \rightarrow g) &= (\tau_{\Box}^i(f' \triangleright \tau_{\Box} g'))^*, \\
t'(f + g) &= \tau_{\Box}^i((\tau_{\Box}(f')^{+D'}) \diamond (\tau_{\Box}(g')^{B'+})), & & \\
t'(g \circ f) &= g' \oplus f'. & &
\end{aligned}$$

LEMMA 5.14. *For every morphism term g , $t(t'(g)) = g$.*

PROOF. By induction on the complexity of g .

LEMMA 5.15. *In each step of our cut elimination procedure, which transforms π into π' , we have $t(\pi) = t(\pi')$ in **BiCartCl**.*

PROOF. Long, tedious but more or less trivial. In steps where we were not precise about the order of application of rules in the transformed proof we rely on some coherence properties, like for example in Case 1.1.3, we use the fact that $t(G(\perp))$ is isomorphic to $\mathbf{O}$ and therefore the order of application of (τ) , (γ) and (θ) is arbitrary.

LEMMA 5.16. *For every morphism term g , there is a mixless proof π of $\mathcal{J}$, such that $g = t(\pi)$.*

PROOF. Let π_1 be $t'(g)$ and let π be the mixless proof obtained from π_1 by our cut elimination procedure. Then by Lemma 5.15, $t(\pi_1) = t(\pi)$, and by Lemma 5.14, $g = t(t'(g)) = t(\pi_1) = t(\pi)$. $\square$

Since the mixless proofs of $\mathcal{J}$ correspond to the constructible canonical transformations, we can derive the following lemma.

LEMMA 5.17. *Every canonical transformation from an arbitrary bicartesian closed category is a subtransformation of a constructible canonical transformation.*

PROOF. By Lemma 5.16 and the universal property of **BiCartCl** it follows that for every canonical transformation α from F to G there exists a constructible canonical transformation β from F to G such that each member of α is equal to a member of β . By our definition, this fact is sufficient for α being a subtransformation of β . $\square$

From Lemmata 3.13 and 5.17 we have the following.

THEOREM 5.4. *Every bicartesian closed canonical transformation is g -dinatural.*

From this theorem and the remark after Example 2.3, it follows that every bicartesian closed canonical transformation is dinatural in the classical sense. Moreover, one has to bear in mind that this property is provable regardless of the choice of language for bicartesian closed categories.

Our proof covers a result from [7] where the authors have used a normalization in a natural deduction system for the fragment of intuitionistic propositional logic that corresponds to cartesian closed categories, to show that all canonical transformations from these categories are dinatural. Since there are still some difficulties with normalization in clumsy λ -calculuses for full intuitionistic

propositional logic, we find an advantage in sequent systems, which are sufficient to deal with the questions of dinaturality. The definitions of operations on objects in the underlying functor category given in [7] are different from our operations in $\mathcal{F}_{\mathcal{B}}$, and a consequence of this difference is that the functor category of [7] is cartesian closed, whereas our $\mathcal{F}_{\mathcal{B}}$ is just symmetric monoidal closed.

Investigations of dinaturality are often tied to investigations of coherence. Some results (cf. [2]) claim that this connection is very strict. However, our graphs, though appropriate for dinaturality, are inadequate for coherence. We leave all this questions about coherence for another occasion.

Acknowledgments. Most of these results are from the author's Ph.D. thesis, written under the direction of Professor Kosta Došen, to whom the author is grateful very much.

References

- [1] Bainbridge, E.S., Freyd, P.J., Scedrov, A., Scott, P.J., 1990, Functorial polymorphism, *Theoretical Computer Science*, 70, pp. 35-64, North-Holland.
- [2] Blute, R., 1993, Linear logic, coherence and dinaturality, *Theoretical Computer Science*, 115, pp. 3-41, Elsevier.
- [3] Došen, K., 1999, *Cut Elimination in Categories*, Trends in Logic – Studia Logica Library, Kluwer Academic Publishers.
- [4] Dubuc, E.J., Street, R., 1970, Dinatural transformations, *Reports of the Midwest Category Seminar*, vol. IV, H. Applegate ed., pp. 126-138, vol. 137, Lecture Notes in Mathematics, Berlin-Heidelberg-New York, Springer.
- [5] Eilenberg, S., and Kelly, G.M., 1966, A Generalization of the Functorial Calculus, *Journal of Algebra* 3, pp. 366-375.
- [6] Gentzen, G., 1935, Untersuchungen über das logische Schließen. *Mathematische Zeitschrift* 39, pp. 176-210, 405-431 (English translation in *The Collected Papers of Gerhard Gentzen*, Szabo, M.E. (ed.), North-Holland).
- [7] Girard, J.-Y., Scedrov, A., and Scott, P.J., 1992, Normal forms and cut-free proofs as natural transformations, in Moschovakis, Y.N., ed., *Logic from Computer Science*, Proceedings of a Workshop held November 13-17, 1989, pp. 217-241, Springer.
- [8] Kelly, G.M., and MacLane, S., 1971, Coherence in Closed Categories, *Journal of Pure and Applied Algebra*, vol. 1, no. 1, pp. 97-140.
- [9] Lambek, J., and Scott, P.J., 1986, *Introduction to Higher Order Categorical Logic*, Cambridge University Press.
- [10] MacLane, S., 1976, Topology and logic as a source of algebra, *Bulletin of the American Mathematical Society*, vol. 82, 1, pp. 1-40.